

Review Article

Microbial Spoilage of Meat and It's Detection - A Review

Nibedita Chowdhury*, Rupsa Sengupta, Saurjaynee Biswas, Deborima Bera, Kakali Bandyopadhyay

*Department of Food Technology, Guru Nanak Institute of Technology,
Panihati, Sodepur, Kolkata, India.*

*Corresponding author's e-mail: nibeditakoyel09@gmail.com

Abstract

Spoilage of meat is a metabolic process which occurs if the meat is left untreated under unsuitable conditions like low oxygen, high water activity & low temperature, hence it becomes unacceptable for human consumption. Meat spoilage is generally caused by unavoidable infections & decomposition of meat by bacteria & fungi. The microorganisms usually come in contact with the meat through the person handling the meat or borne in the animal itself. Unattractive odours & flavours, discolouration, gas & slime are produced by the microbes associated with deterioration of meat. For so long, food industry has been looking for technologies to detect deterioration of meat. Few detection methods are being used in food industry these days to ensure freshness of meat & customer satisfaction such as spectroscopy, use of AI methods, biosensors etc. In the present review article, it has been focused on the detection of meat spoilage by FT-IR spectroscopy with PLS Analysis method, Xanthine Oxidase enzyme biosensor method & Enumeration method.

Keywords: Spoilage of meat; Water activity; Spectroscopy; AI methods; Biosensors; Xanthine; Oxidase enzyme; Enumeration Method.

Introduction

The demand for poultry and poultry products is increasing day by day in both international and domestic markets [1] in order to protect and maintain human-health. Safety and quality are considered as the most important concerns in the food industry as they are directly related to get healthy life. Our human body requires proper and maintained diet comprising of all required nutrition to carry on the daily life-functions [2].

This required diet contains macro - nutrients and micro nutrients which can be found from a number of food source including cereals, meat, legumes, fruits, milk and vegetables [3]. Being nutritious, meat provides energy to the human body to perform daily activities by fulfilling the requirements needed by the body [4]. Meat is considered as a rich source of amino acids, essential fatty acids and vitamins [5].

Meat is a good source of nutrition and different kinds of meat (e.g., chicken mutton, lamb, beef, fish etc.) has different composition with different nutritional values [6]. Due to increase in concerns of consumer over food quality and food safety, it becomes a very

important need for Industries to ensure safe, unspoiled and healthy food to the consumer as food is related to increase in morbidity, mortality, human suffering and also affects economy [7]. To comply with the need a momentous effort made by meat industry has been observed to ensure the quality and safety of meat products which includes food spoilage, food poisoning and food waste [8].

Meat gets spoiled for various reasons but microbiological spoilage in meat occurs with the growth of microorganisms. The spoilage of meat is identified by its organoleptic changes which make the meat unacceptable to eat [9]. These changes include appearance (discoloration), changes in odours, changes in taste, slime formation. This may result from formation of metabolites, decompositions, conversion of organic and inorganic matter and by enzymatic activities which occur within tissues after post-mortem can lead to change during storage time due to the growth of microorganisms [9]. Basically, food gets rejected when it gets spoiled by triggering some characteristics which makes the meat unacceptable to the consumer [10].



Detection Methods of Non-permitted Food Color, Metanil Yellow: A Review

Parag Chattopadhyay¹ | Aditya Paul¹ | Samridhya Paul¹ | Sukanya Ray¹ | Kakali Bandyopadhyay²

¹Department of Food Technology, Guru Nanak Institute of Technology, Kolkata, India.

²H.O.D and Associate Professor, Department of Food Technology, Guru Nanak Institute of Technology, Kolkata, India.

To Cite this Article

Parag Chattopadhyay, Aditya Paul, Samridhya Paul, Sukanya Ray and Kakali Bandyopadhyay, "Detection Methods of Non-permitted Food Color, Metanil Yellow: A Review", *International Journal for Modern Trends in Science and Technology*, Vol. 07, Issue 03, March 2021, pp.: 201-204.

Article Info

Received on 15-February-2021, Revised on 12-March-2021, Accepted on 17-March-2021, Published on 20-March-2021.

ABSTRACT

Food adulteration is a primary global concern for public health, especially in developing countries, due to the lack of monitoring and appropriate policy developments and executions. Due to its high demand in international trade, turmeric (*Curcuma Longa*) is subjected to economically motivated chemically unsafe adulteration, namely metanil yellow. Metanil yellow (3-4)-anilinophenylazo) benzene sulfonic acid sodium salt), is a hazardous dye and a common adulterant used in turmeric powder and other yellow colored food products. The toxic chemical travels in blood and reaches various organs and interferes with various cellular metabolic processes there. Our earlier studies reveal that metanil yellow generates oxidative stress in various vital organs such as heart, liver, and kidneys. As per the guidelines of food quality some conventional methods are used but these techniques possess various limitations. This study aims to review the use of FT-Raman and FT-IR spectroscopy for evaluation of metanil yellow in turmeric powder.

KEYWORDS: Non-permitted food colors, metanil yellow, turmeric powder

INTRODUCTION

There are several types of dyes used as food additives and food colorants. Metanil yellow is a yellow dye used extensively as a food colorant. It is made from diazotized metanilic acid and diphenylamine [1]. Azo dyes are also used in laboratories as biological indicators. Turmeric (*Curcuma long L.*) is an appetizing root commonly used for food seasoning and for medicinal purposes. Turmeric has a long history of medicinal use in Asian countries (Reema F Tayyem et al. *Nutr Cancer*. 2006[2]) and is used in root, oil, and powder forms. Its medicinal value is mainly due to its content of curcumin (diferuloyol methane) (T Osawa et al. *Biosci Biotechnol Biochem*. 1995 Sep [3]) with attributed medical properties including anti-inflammatory, anticarcinogenic,

antioxidant, and wound-healing effects (B Joe et al. *Crit Rev Food Sci Nutr*. 2004[4]). Curcumin has also been reported to have promise for development of therapies for Alzheimer's disease (Tsuyoshi Hamaguchi et al. *CNS Neurosci Ther*. 2010 Oct [5]).

According to the sources it has shown that the nutrient and acidity content in soil, fertilizer, soil type and cultivar affects the curcumin content in turmeric. Reported curcumin concentrations in turmeric range from 0.3% to 8.6% [3,6-9]. Curcumin is isolated from turmeric for medicinal and cosmetic purposes. Although whole, dried, or fresh turmeric are mostly free of contamination, turmeric powder can be deteriorated with different chemical powders used as substitutes for curcumin (Sasikumar.B et al. 2004 [10]). It has been using in many unorganized



Synthesis of activated carbon material using sawdust as precursor and its application for dye removal: batch study and optimization using response surface methodology

Dolanchapa Sikdar^{1,2} · Sudipta Goswami¹ · Papita Das¹

Received: 17 December 2020 / Revised: 5 February 2021 / Accepted: 15 February 2021

© The Author(s), under exclusive licence to Springer-Verlag GmbH, DE part of Springer Nature 2021

Abstract

In the present study, an inexpensive adsorbent has been synthesized from waste sawdust which can remove the highly toxic dye indigo carmine (IC) from aqueous solution. Activated carbon from sawdust was synthesized using physical and chemical treatment. After using the adsorbent in batch-dye adsorption experiments, it has been characterized by SEM, FT-IR and XRD which showed significant up-taking capacity (77%) of the dye on its surface. Sensitivity of the dye removal process has been analysed by undergoing batch study with varying initial concentrations of the IC dye, adsorbent dose, pH and temperature and optimization of the process which indicated the optimum condition of the experimental parameters. Various isotherm models and kinetic models have been used for fitting of experimental data and examined to analyse efficiency of the adsorption mechanism. Langmuir isotherm and pseudo-second-order kinetics were found to be followed by adsorption process over the entire dataset of tested experimental results. The maximum uptake capacity was found to be 30 mg g^{-1} . Thermodynamic investigation was also conducted to project the spontaneity of the reaction.

Keywords Sawdust · Waste biomass · Biochar · Low-cost adsorbent · Dye removal · Indigo carmine (IC) · Adsorption · Environmental pollution

1 Introduction

Various industries (paper, leather, food, cosmetic, textile, etc.) use dye for colouration of their product [1, 2] and thus ending up discharging a considerable amount of dye in water body in the form of waste water. Discharge of waste water containing dye discharges into the environment causes detrimental effect on flora, fauna and human beings [3, 4]. There are various methods (physical, chemical, biological) that are being employed to remove excess dye from the waste water before discharging into the environment [5, 6]. Among all other techniques, the superior one is found to be the adsorption technique with respect to low

cost, ease of operation and simplicity of design [7, 8]. The most widely used adsorbent is activated carbon because of high pore size and surface area [5].

Indigo carmine (IC) is a synthetic blue colourant (acid dye) and comprises of two sulfonated groups (negatively charged) coupled with four benzene rings. It is a typical recalcitrant dye with high solubility (10 g/l) [9] and used as colouring agent in various industries. Discharge of IC dye beyond permissible limit can cause potential risk to human health and cause development of acute neurotoxicity [10], hypertension and cardiovascular disease [11, 12], diarrhoea, nausea etc. [13, 14]. It is a highly toxic dye and even touching may cause human eye and skin irritation and may cause permanent damage to cornea [15, 16].

As taking in consideration of the toxicity of the dye, diverse conventional techniques are used like photocatalytic degradation, electrodialysis, coagulation, ozonation and adsorption for the removal of IC dye from polluted water [17–21]. Though these processes are tedious and non-economical, however, the adsorption technique among all these techniques is found to have shown high removal efficiency and better operational characteristics [22].

✉ Dolanchapa Sikdar
dolanchapasikdar88@gmail.com

¹ Department of Chemical Engineering, Jadavpur University, Kolkata 700032, India

² Department of Food Technology, Guru Nanak Institute of Technology, Kolkata 700114, India



A Review on Cocoa Butter Alternatives in Chocolate Preparation

Writtika Das¹ | Sweta Das² | Shairee Ganguly³ | Dolanchapa Sikdar⁴ | Kakali Bandyopadhyay⁵

^{1,2}: B. Tech. Students,

^{3,4}: Assistant Professor

⁵: Associate Professor and HOD

Department of Food Technology, Guru Nanak Institute of Technology, 157/F, Nilgunj Road, Sodepur, Panihati, Kolkata 700114

To Cite this Article

Writtika Das, Sweta Das, Shairee Ganguly, Dolanchapa Sikdar and Kakali Bandyopadhyay, "A Review on Cocoa Butter Alternatives in Chocolate Preparation", *International Journal for Modern Trends in Science and Technology*, Vol. 07, Issue 01, January 2021, pp.- 15-140.

Article Info

Received on 11-December-2020, Revised on 30-December-2020, Accepted on 06-January-2021, Published on 19-January-2021.

ABSTRACT

Cocoa butter naturally occurs in cocoa bean. It is present about 50% of cocoa nib. It is highly resistant to oxidation due to presence of high level of natural tocopherol. It is brittle at room temperature and its melting point is between 34° C to 38° C. But, cocoa butter is expensive and its price is subjected to large fluctuations. That's why some alternatives for cocoa butters are produced, such as cocoa – butter equivalents (CBEs, like from enzymatic inter-esterification of tea seed oil and fatty acid methyl esters), cocoa-butter substitute (CBSs, esterifies propoxylated glycerin containing acyl groups derived from saturated linear fatty acids, at least 50 moles percent of the total acyl groups are used) and cocoa-butter replacers (CBRs, like evaluation of milk fat fractional and modified techniques for making CBR.) Their composition according to triglycerides, fatty acids, sterols and other unsaponifiable components are discussed in this paper. Coconut oil, non-lauric contained fats like palm oil, soybean oil, rapeseed oil, can be used as replacer. These alternatives have various advantages; it improves fat stability, reduces fat migration, and incorporates softness to the product. As the alternatives do not require tempering, it is easier to achieve glassy texture. This work reviews on the theory of the compositional data of vegetable oils, and fats which are used as cocoa – butter alternatives in the production of chocolate.

KEYWORDS: cocoa butter alternatives, melting point, chocolate

I. INTRODUCTION

Cocoa bean (CB) is the fatty seed found inside a cocoa pod, fruit of the

Theobroma cocoa plant. It is a small evergreen tree that belongs to the family Malvaceae. There are several processes which undergo the preparation of cocoa butter. It is brittle in Temperature below 25°C soften in hand and melts in mouth having a

temperature of about 34°C [33]. Cocoa butter has several advantages which makes it the main ingredient saturated fat and less amount of caffeine and theobromine. It also contain some vitamins in the soluble form such as vitamin E in the form of beta – tocopherol, alpha – tocopherol and gamma – tocopherol. Among all this form of tocopherol, only beta – tocopherol is used for

Organizing The Unorganized: An Overview of COVID-19 Impact On FMCG Sector and Indian Economy

Saikat Mazumder^{1*}, Kakali Bandyopadhyay², Rajdeep Chakraborty³, Ayan Malakar³, Parna Sarkar³

^{1*} Assistant Professor, ² Head, Associate Professor, ³ B. Tech. Students
Department of Food Technology, Guru Nanak Institute of Technology

Abstract

Fast-moving consumer goods or FMCG sectors have a significant role in the Indian economy. This part is now confronting some issues because of the significant lockdown in the light of the new corona virus which resulted in the occurrence of the pandemic. There are two market divisions of FMCG sectors, they're organized and unorganized, uniquely the chaotic part is unorganized sectors are greatly influenced by the lockdown in pandemonium. The FMCG sectors have indicated an amazing development for recent years. The aim of the review to focus the impact of a worldwide pandemic on the FMCG segment in India and the behavioral changes of customer and how the FMCG sector will boost the economy. The government polices and different technology stack like automation that can be included in the FMCG sector can bounce back and boost the economy. Furthermore, more investment particularly on livestock sector can contribute 1% to the annual GDP. Alongside, a continuous effort should make in public private partnership model to create bridge between the organized and unorganized sector.

Keywords: FMCG, Covid-19, Food Sector, Technology stack, organized sector, unorganized sector, public private partnership

1. Introduction

India is the 5th largest economic force in the world with 3.202 trillion US\$ (Nominal; 2020). FMCG is the 4th biggest economy supplier in India with a market size of 52.75 billion US\$. Fast-moving consumer goods are non-durable products that sell quickly at relatively low cost. FMCGs have low-profit margins Fast-moving consumer goods are an inexpensive product. These items are perishable and require small shopping endeavors. FMCG sectors have a wide assortment of the range including biscuit packets to toiletries. This division has provided very nearly 3 million work openings, which is nearly 5% of the total factory employment (Dabur annual report; 2017), but has seen a huge decline in the current scenario but interestingly the food market has seen a surge of almost 40%. Most of the essential food items are out of stock in the market because the majority of the food sectors are unorganized market player. There are two different divisions in FMCG sectors, they're organized and unorganized. But these unorganized parts are facing major problems due to the pandemic situation which results in the lockdown of markets. The majority of the absolute workforce of the nation is from the unorganized sectors. The organized sectors have confronted this equivalent circumstance yet but they're bouncing back rapidly. We'll discuss in this paper the behavioral changes of consumers in respect of FMCG sectors in this pandemic situation and some advisory solutions to overcome this uncertainty of unorganized sectors and to bring the unorganized market played into an organized market player. While the unorganized market player which runs our informal income are the ones which are in higher risk in this pandemic and it's expected that more than 400 million people which help running our informal economy may lose their job.

2. IMPACT OF COVID 19 ON FOOD SECTOR (FMCG) AND LIVELIHOOD IN INDIA

With the intent to stop the spread of viruses the lockdown has been imposed and the situation remains volatile with the trajectory of the corona virus undetermined and a significant fall in the economic fallout according to HUL chairman - Sanjeev Mehta [1]. The government views the pattern of the spread of COVID-19 as similar to the 2009 H1N1 influenza pandemic, meaning the spread is unlikely to be uniform. After the unlock III, it is quite expected to maintain the full lockdown in "hotspot" areas and relax it in other places and started to unlock the country once again with full flow but with safety measures. While in some places across the Country and various states, the state government decided to impose strict lockdown again due to the current scenario of COVID-19. The FMCG sector has seen a huge decline in the current scenario but interestingly the food market has seen surge online sales were a mere 2.8% of the overall sales for FMCG companies in the first quarter of 2020. Kerala-based packaged foods company Elite Foods saw a 100% jump in online sales in certain regions for its products that include cakes, bread and buns, Indian instant mixes,

A Review on Utilization of Biosensors for Detection of Adulteration in Fish

Dr. Kakali Bandyopadhyay¹ | Soumily Misra² | Somdeepa Bhattacharya² | Arnab Mukherjee² | Arunanshu Shee²

¹ H.O.D and Associate Professor, Department of Food technology, Guru Nanak Institute of Technology, Kolkata, India.

² UG Students, Department of Food technology, Guru Nanak Institute of Technology, Kolkata, India.

To Cite this Article

Dr. Kakali Bandyopadhyay, Soumily Misra, Somdeepa Bhattacharya, Arnab Mukherjee and Arunanshu Shee, "A Review on Utilization of Biosensors for Detection of Adulteration in Fish", *International Journal for Modern Trends in Science and Technology*, 6(10): 50-55, 2020.

Article Info

Received on 12-September-2020, Revised on 28-September-2020, Accepted on 04-October-2020, Published on 08-October-2020.

ABSTRACT

The consumption of fish has lately seen an upward trend in recent days considering the wide range of health benefits of fish and sea foods and with it, the issues of prevalent adulteration of fresh fish with unapproved chemicals and additives have come into light both through electronic and print media. Several instances of adulteration of fish and fishery products with unsafe chemicals like formaldehyde, ammonia and additives like sodium benzoate has been highlighted in a number of reports and research papers. Presence of these adulterants in freshly marketed fish seriously compromises health of consumers and daily ingestion of considerable amounts of these adulterants can generally cause severe abdominal pain, coma, renal injury, vomiting and possible death. This has aroused concerns regarding the safety and quality of fish as well as the necessity of a reliable and faster technology which can easily identify the contaminants and toxicants in food materials. As per regulatory guidelines of food quality some conventional methods are used but these conventional techniques possess various limitations. Biosensor is an analytical device which can replace these conventional techniques and offers a quick on-site monitoring with accuracy and provides an index of quality of the product in real time. This study aims to review the recent advancement in biosensors and their contributions in determination of adulteration and freshness concerning fish and seafood safety.

KEYWORDS: Adulteration of fish, Formaldehyde, Ammonia, sodium benzoate, Biosensor.

INTRODUCTION

Fish is a healthy food and serves as a most important source of animal proteins and great source of ω -3 PUFAs and micronutrients for people worldwide (Mohanty et al. [1]) mainly in the underdeveloped and developing countries. High consumer demand for fish and fish products has resulted in enhanced fish trade in various countries. To cater the need of consumers, fish is exported from the place of production to various distances. As fish is perishable, the flesh of fish can spoil quickly and it should be eaten on the day of capture, unless cured (Marzuki et al. [2]). In order

to extend the storage life as well as to improve the sensory attributes i.e. appearance, fishermen and fish vendors tend to carelessly use hazardous chemicals like formaldehyde and ammonia and additives such as sodium benzoate in various fish and sea foods. Formaldehyde has been identified "as carcinogenic to human" by IARC (International Agency for Research on Cancer) (Bianchi [3]). As consumption of these adulterants is hazardous to human body, it is important to investigate the presence of adulterants as well as the quality of fish. Various methods have been adopted for detection of adulterants like liquid



STUDY ON THE FORTIFICATION TECHNIQUES OF MULTIPLE FOOD PRODUCTS USING IMMUNITY BOOSTING MICRONUTRIENTS AND ANTIOXIDANTS: A LITERATURE REVIEW

Kakali Bandyopadhyay
H.O.D and Associate Professor
Department of Food technology
Guru Nanak Institute of Technology
Kolkata, India.

Mariya Mukherjee, Pulak Deb, Akash Biswas, Riya Kumari Gope, Sweta Das
UG Students
Department of Food technology
Guru Nanak Institute of Technology
Kolkata, India

Abstract: During this ongoing period of the devastating pandemic of COVID-19 caused by Novel Coronavirus, the term “Immunity” has become quite a catchword, and a strong and healthy immune system has become the need of the hour to build up our body resistance against pathogens and provide protection against them. Researches on various nutritional components and their bioavailability point to the fact that there are several micronutrients and trace elements whose intake through diet can help in triggering up the functioning of the immune system. Nowadays, as per the Food Safety and Standards (Fortification of Foods) Regulations, it is also advised that we must buy and consume products that have the “+F” logo or fortification logo which means that those products are enriched in certain added vitamins and trace metals from which the benefits of a stronger immune response can be obtained. It has been observed that among the vitamins, vitamin A, C, D and E and the mineral zinc play a significant role in boosting up the immune response through their interactions with the cells of the immune system. Also, antioxidants like β -carotene, anthocyanins, etc play a major role in the immune system functioning by lowering down the level of oxidative stresses in the body. These immunity boosting vitamins, minerals and antioxidants can be incorporated into regular food products through the method of fortification of the products or even, they can be added into an immunity boosting drink, powder or supplements. Various researches and reviews demonstrate how these micronutrients can be added or incorporated into a food product; they can be obtained from artificial/synthetic sources or natural sources like fruits containing these

vitamins and minerals can be used as functional ingredients in the products. Sometimes, multivitamins or minerals can be incorporated into the same product. Therefore, the following review will be based on the study of the methods in which various products have been fortified from time to time with immunity boosting nutrients like vitamins A, C, D and E, zinc and antioxidants as demonstrated in numerous research work. It will be a detailed study on the techniques followed during the fortification procedures as well as the sources from which those micronutrients and antioxidants are obtained. The roles played by the micronutrients in building immunity will also be discussed.

Index Terms - Immunity, Fortification, Bioavailability, Vitamins, Micronutrients, Antioxidants, Minerals, Regular food products

I. INTRODUCTION

Immunity is the body’s internal capability to resist the harmful effects of an invading pathogenic microorganism or even minimize the effects of infections caused by them. It is mostly an internal factor of the body and the strength of a person’s immunity depends mainly on the strength of the immune response generated by his immune cells. However, there are certain micronutrients which are observed to enhance this overall functioning of the immune system and these elements are mostly obtained from consuming foods which are rich in them. The vitamins A, C, D, E and the mineral zinc are the micronutrients which are observed to perform such immunity boosting activities. Each of these micronutrients has its own distinct mechanism through which it triggers the

Onion Productivity and Price Change Aspects in India: An Overview

Supriti Manna¹ | Dabasmitta Saha¹ | Subham Roy Chowdhury¹ | Amit Barman¹ | Saikat Mazumder¹

¹Department of Food Technology, Guru Nanak Institute of Technology, Kolkata, West Bengal, India

To Cite this Article

Supriti Manna, Dabasmitta Saha, Subham Roy Chowdhury, Amit Barman and Saikat Mazumder, "Onion Productivity and Price Change Aspects in India: An Overview", *International Journal for Modern Trends in Science and Technology*, 6(10): 118-122, 2020.

Article Info

Received on 30-September-2020, Revised on 12-October-2020, Accepted on 15-October-2020, Published on 23-October-2020.

ABSTRACT

Allium Cepath scientific name of an onion. After potato Onion is the second most essential vegetable in our India. And also onions are very famous in all around the world. In India, the productivity of onion is about 13 lakh tons every month, and in the world, the onion productivity ratio is about 19.40 million. There are about 100 types of onion throughout the world and basically, India produces 9 types of major onion. Onion is very rich in vitamin C (12%) content and vitamin B-6 (5%) content and also rich in minerals like iron (1%) and calcium (2%). In the last 10 years, Maharashtra (4905.0 thousand tons) is the biggest onion producing state in India. But in 2019 onion prices overlap at Rs 10 kg from Rs 100 kg as demand loss will take place. A huge amount of wastage of onion takes place every year due to bad Monsoon, transport system and the amount of wastage is nearly 25% of the total production. In 10 years a remarkable increase in the production of onion in India and also an increase in the area about 768,000 ha to 1064,000 ha. Onion production has been several benefits like it reduces our cholesterol level, fights against inflammation, decreases triglycerides and it contains a high amount of antioxidants. The impact of onion production on the Indian economy, an increase of onion productivity, price drop, Government policies related to onion production also have been focused on in this review.

KEYWORDS: Onion production, government export policies, Wholesale price, Price change, Storage of onion

I. INTRODUCTION

Onion is the second largest production crop in India and also all over the world [1]. The best season for onion production in India is August October, and for hilly areas, the best time for onion production is September to December [2].

Throughout the world, China is the most onion producing country and produces 20507.76MT onions whereas India produced 15118.00MT onions per year. Maharashtra is the best onion producing state in India, produced 8474.73 thousand tons of onion per year. But As per the current report onion is expected to be 20% less in

the country and also a reduction of 42% expected in Maharashtra due to bad climate conditions [3].

The average wholesale cost of onion in India is 1280.32 Rs/quintal. And a total of 15.89 lakh MT Onion is exported in 2018 and in 2019 the export structure is 15.22 lakh MT [4,7,9]

Besides production argumentation, there are several policy tools available to arrest the increasing prices such as the withdrawal Of Merchandise Export Incentive scheme (MEIS) the rewards under MEIS mostly in form of transferable duty credit scripts and to encourage exports and curb free fall in prices for Onion at home, the MEIS

RESEARCH

Open Access



Activated carbonaceous materials from tea waste and its removal capacity of indigo carmine present in solution: synthesis, batch and optimization study

Dolanchapa Sikdar^{1,2*}, Sudipta Goswami¹ and Papita Das¹

Abstract

The present work determines efficiency of domestic food waste like tea waste in removing indigo carmine from synthetic textile wastewater. Carbonaceous material (tea waste) has been employed and it showed removal percentage of indigo carmine at 90% through adsorption process. Carbonaceous material was chemically activated by using modified Hummer's method and it was observed through Scanning Electron Microscopic image, Fourier Transform Infrared Spectrometer, X-ray diffractometer and Brunauer, Emmett and Teller analysis that revealed that the modified biochar is comparably similar to graphene oxide (GO). Various experimental parameters are evaluated for the removal efficiency of the synthesized adsorbent under the present study. Results of the experiments performed using the GO-like adsorbent synthesized from tea waste confirmed potential efficiency of adsorption of indigo carmine dye from synthetic waste water solution. The adsorption mechanism has been analysed by fitting the experimental data in different adsorption isotherm and kinetic models. The results indicated that the adsorption followed Langmuir isotherm model with maximum uptake of 20 mg g^{-1} and pseudo second order kinetic model with the best correlation coefficient. The thermodynamic study showed the dye removal to be spontaneous and endothermic.

Keywords: Indigo carmine (IC), Adsorption, Tea waste, Environmental pollution

Introduction

Dyes are a specific group of chemical pollutants, posing serious environmental problems, due to annual production in large amounts from different industrial sector like textile industry [1–3]. If this wastewater is released into a natural aquatic system, the photochemical activities of that aquatic system are adversely affected by the coloration of dye which reduces light penetration. Carcinogenic dyes might cause mutations in organisms [4, 5]. Dyes are chemically designed and manufactured to be resistant to degradation by

oxidizing agents, light, high temperatures and hence, cannot be easily removed by the conventional wastewater treatments methods [6, 7].

Indigo carmine (IC) dye is widely present in wastewater produced from textile, food, cosmetic, pharmaceutical and paper industries [8–12]. According to WHO, the permissible limit of indigo carmine dye is $5 \mu\text{g L}^{-1}$ in water [13]. Wastewater containing IC dye must undergo treatment before discharging into the environment as it may cause skin and eye irritation, cancer to human [14–16]. Various methods that are based on aerobic or anaerobic biodegradation and photodegradation are not 100% efficient for removal of most of the dyes from wastewater. The various

* Correspondence: dolanchapasikdar88@gmail.com

¹Department of Chemical Engineering, Jadavpur University, Kolkata 700032, India

²Department of Food Technology, Guru Nanak Institute of Technology, Kolkata 700114, India



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Overview of Worldwide Potato Production Scenario with Special Focus in India and Its Industrial Utilization

Debdatta Sarkar¹ | Bisakha Saha¹ | Arunima Singha¹ | Saikat Mazumder¹

¹Department of Food Technology, Guru Nanak Institute of Technology, Kolkata, West Bengal, India

To Cite this Article

Debdatta Sarkar, Bisakha Saha, Arunima Singha and Saikat Mazumder, "Overview of Worldwide Potato Production Scenario with Special Focus in India and Its Industrial Utilization", *International Journal for Modern Trends in Science and Technology*, 6(11): 137-142, 2020.

Article Info

Received on 26-October-2020, Revised on 10-November-2020, Accepted on 20-November-2020, Published on 23-November-2020.

ABSTRACT

India became the planet's second-largest potato producing country all over the world because potato output increased from 1.3 million to over 34 million over the last six decades. Despite that, this spectacular increase makes a series of less readily apparent tendencies in the growth rates for potato production, area, and yields. When the area harvested enlarged, the growth for area harvested touched a series of peaks and valleys; as yields per hectare continuously increased, the growth rate for productivity gradually decreased to a half. In the present year, Uttar Pradesh is the major Potato producing State with 31.26% of production share, followed by West Bengal, Bihar, Gujarat, and Madhya Pradesh with 23.29 %, 13.22%, 7.43%, and 6.20% share respectively. This paper analysis is based on the driving factors behind the increase in production, area harvested, and yields. Sub-sector specific recommendations include greater eco-efficiency in cold storage.

KEYWORDS: Potato production, potato Nutrition, Potato varieties, Industrial utilization

I. INTRODUCTION

Potato is a root vegetable along with that it is also a cool-season vegetable which is in part with wheat and rice because it is one of the most important staple crops in the human diet all around the world. Potato is a specialized underground storage stem which is also known as "tuber". According to FAO (the Food and Agriculture Organization) potato is devoured by more than one billion people around the world. It is a high-quality vegetable as well as food crop used in the reparation of more than 100 types of recipes in INDIA. The protein of potato has high biological value than protein in cereals and milk [1].

Potatoes are rich in vitamins, minerals, and

antioxidants, which make them very healthy. Studies say that potatoes have a variety of impressive health benefits, including improved blood sugar control, reduced heart disease, etc. One medium baked potato (173 grams) including the skin, provides Calories: 161 grams Fat: 0.2 grams, Protein: 4.3 grams Carbs: 36.6 grams Fibre: 3.8 grams Vitamin C: 28% of the RDI Vitamin B6: 27% of the RDI Potassium: 26% of the RDI Manganese: 19% of the RDI Magnesium: 12% of the RDI Phosphorus: 12% of the RDI Niacin: 12% of the RDI Folate: 12% of the RDI The nutritional content of potatoes can vary by the variety of it and how they are prepared. For example, frying potatoes adds more calories and fat than baked

A Review on Anthocyanin Pigments with respect to its Nutraceutical Properties

Srijeeta Saha¹ | Shairee Ganguly¹ | Dolanchapa Sikdar¹

¹ Department of Food Technology, GURU NANAK INSTITUTE OF TECHNOLOGY, KOLKATA, West Bengal, India

To Cite this Article

Srijeeta Saha, Shairee Ganguly and Dolanchapa Sikdar, "A Review on Anthocyanin Pigments with respect to its Nutraceutical Properties", *International Journal for Modern Trends in Science and Technology*, 6(12): 54-60, 2020.

Article Info

Received on 07-November-2020, Revised on 22-November-2020, Accepted on 28-November-2020, Published on 01-December-2020.

ABSTRACT

Anthocyanin pigments are readily degraded during processing and storage of foodstuffs that can have a dramatic impact on color quality and may also affect the nutritional properties. Total anthocyanin pigment content and indices for polymeric color and browning are easily measured with simple spectrophotometric methods. Once individual pigments are identified, their changes can be monitored by high-performance liquid chromatography (HPLC). The edible fruits of 12 plants were extracted in methanol and subjected to solvent-solvent partitioning to yield three fractions, hexane, ethyl acetate, and aqueous. A number of factors affecting anthocyanin stability and color are discussed in this review. Anthocyanins are probably the most spectacular of plant pigments since they are responsible for most of the red, purple and blue pigmentation of flowers, fruits and vegetables. However, because of their highly reactive nature, anthocyanins readily degrade, or react with other constituents in the media, to form colorless or brown colored compounds. The presence of an oxonium ion adjacent to carbon 2 makes the anthocyanins particularly susceptible to nucleophilic attack by such compounds as sulfur dioxide, ascorbic acid, hydrogen peroxide and even water. Loss of anthocyanin pigmentation also occurs in the presence of oxygen and various enzymes, and as a result of high temperature processing. Certain degree of pigment stabilization may be conferred by acylation with various organic acids, co pigmentation, self-association and/or metal chelation. In addition, pH has a marked effect on anthocyanin stability, and on the color of media containing these pigments. A number of anthocyanin-rich sources have been investigated for their potential as commercial pigment extracts. Although their application is primarily limited to acidic media, continued research on the chemistry of anthocyanins may lead to application and stabilization of these pigments in a wider variety of food products.

KEYWORDS: Anthocyanins, Natural pigment, Pigmentation, acylation, chelation, stabilization, fortification.

I. INTRODUCTION

Anthocyanins are water-soluble vacuolar pigments which depend on their pH, which may appear red, purple, blue or black. There are many food plants which are rich in anthocyanins include the blueberry, raspberry, black rice, and black soybean and many others pigments colored red, blue, purple or black.^[12] Anthocyanins comprise a

group of naturally occurring pigments which are responsible for the blue, red, purple, violet and magenta coloration of most species in the plant kingdom. These polyphenolic substances are glycosides of anthocyanins, polyhydroxy and polymethoxy derivatives of 2-phenylbenzopyrylium or flavylum salts. The large number of glycosyl and acyl groups which may bind to the sixteen different naturally occurring anthocyanidins has



HIBISCUS PETAL EXTRACTION FORTIFIED FRUIT JUICE- A HEALTH DRINK.

AUTHORS: - KAKALI BANDYPADHYAY¹, SHAIREE GANGULY², PRITHWEEJIT BHATTACHARYYA³, LOPAMUDRA BANERJEE⁴

*1: HEAD AND ASSOCIATE PROFESSOR

2: ASSISTANT PROFESSOR

3 and 4: B.TECH STUDENTS

1, 2, 3 AND 4: DEPARTMENT OF FOOD TECHNOLOGY, GURU NANAK INSTITUTE OF TECHNOLOGY, INDIA, EMAIL:

Abstract:-

The aim of this paper was to bring out the beneficial properties present in petals of hibiscus flowers and their fortification in fruit juice. One of the flowers of Indian Subcontinent is "Hibiscus rosa sinensis" which is rich in antioxidants, anthocyanins, flavonoids. Main coloring pigment is anthocyanin which is proved to cure cancer, lower down cholesterol level and clear the digestive tract. Freshly plucked hibiscus flowers were taken and their petals were removed by hand. Petals were put in a tray drier for drying at 51.8°C. After drying they were crushed into fine pieces manually. 100 grams of powdered leaves were obtained. Anthocyanin being water soluble, the powder was boiled in water at 80°C. The anthocyanin pigment was extracted in liquid form. It was mixed with lime juice and polyphenol content was measured by gallic acid equivalent method. Final product obtained showed 13 times increase in polyphenol content compared to commercially available fruit juice. Hence the above juice can be used as an effective health drink for human consumption and protect the body from free radical damage.

Keywords: - Anthocyanins, polyphenols, gallic acid equivalent method.

INTRODUCTION

Flowers and other parts of the plant are used to make medicine. Hibiscus is used for conditions such as high blood pressure, high cholesterol, increasing the production of breast milk etc. It is a flower which is widely grown in the Indian subcontinent and is available around the year. Many varieties are available worldwide among which the variety which is widely available in India is "Hibiscus rosa sinensis". The herb "Hibiscus rosa sinensis" (Malvaceae) is native to China. Many species of Hibiscus are grown for their showy flowers. It is a shrub widely cultivated in the tropics as an ornamental plant and has several forms with varying colors of flowers. Hibiscus has also medicinal properties and takes part as a primary ingredient in many herbal teas. The red flowered variety is preferred in medicine as reported by Vincenta Khristi and V. H. Patel (2nd November, 2016). According to Pragya Singh et al (31st May, 2017), the flower plays a crucial role in treating medical problems including many cardiovascular disorders, helmenthic disease and cancer. The plant also act as an anti oxidant and used in obesity management. The vitamin C naturally found within the hibiscus works to complement the iron, increasing its absorption in blood and acts effective cure for anemic patients. Anthocyanin-rich mixtures of bioflavonoids may provide protection from DNA cleavage, estrogenic activity, enzyme inhibition, boosting production of cytokines, anti-inflammatory activity, decreasing capillary permeability and fragility, and membrane strengthening Mary Ann Lila (12th May, 2004). Being water soluble the pigment can be easily extracted by using water and thus by fortifying it with regular food items can serve as beneficial and cheap source of nutritional super foods. Studies of Yasuyuki Nakamura et al (8th September, 2014) showed that red colored pigments which are present in the flower are mainly anthocyanins which have been widely used as coloring agents. Major content present is cyanidin-3- sambubioside and delphinidin-3-sambubioside from which the conclusion comes that the major component in the petals of hibiscus flower is cyaniding-3-sambubioside, although it contain many other strains of anthocyanins including many hybrid forms so it is still a question that whether every hibiscus flower produces cyaniding-3-sambubioside as the major anthocyanin flora or not. Research of Jadhav et al (2nd July, 2009) all the parts of "Hibiscus rosa sinensis" and chemical constituents are used as anti-tumor, antifertility, antiovolutory, ant implantation, anti-inflammatory, analgesic, ant estrogenic, antipyretic, antispasmodic, antiviral, antifungal, antibacterial, hypoglycaemic, spasmolytic, CNS depressant, hypotensive and juvenoid activity. As recommended by the "World Health Organization" the traditional health and folk medicine systems have proved to be more effective in health problems worldwide. Hibiscus flower is certain to emerge in the near future as a major player in the growing field of herbal health supplements and medicines both in daily self-care and in professionally managed health care system.

Development and Characterization of Biocolour Fortified Yogurt: A New Pathway towards Functional Foods

Shairee Ganguly^{#1}, Kakali Bandyopadhyay^{#2}, Riya Dasgupta^{*3}

^{#1}Assistant Professor, ^{#2}Associate Professor and HOD,

[#]Department of Food Technology, Guru Nanak Institute of Technology, 157/F, Nilgunj Road, Sodepur, Panihati, Kolkata 70011, West Bengal, India.

^{*3}Student, Department of Chemical Technology, University of Calcutta, A.P.C. Road, Kolkata, West Bengal, India

Abstract: At present, the demand for functional foods is increasing worldwide due to the increased awareness on therapeutic and medicinal properties and their benefits among public because they provide health benefits beyond the provision of essential nutrients. Numerous plant foods or physiologically active ingredients derived from plants have been investigated for their role in disease prevention and health. Natural dyes are those derived from plants, insects, animals and minerals. In this study, color extracted from purple cabbage was used for the development of fortified yogurt, since dairy products or the milk derived products can be considered as functional foods as they are health beneficial for human. The physicochemical and rheological properties of fortified and plain yogurt were checked. Plain yogurt is considered as control. The rheological properties were measured using rheometer which shows a similar flow behavior of Bingham plastic nature. The color stability in terms of anthocyanin content was determined at 560nm at different pH. The polyphenol content (in terms of mg GAE/ gm dry sample) of both fortified and control yogurt were determined spectrophotometrically which shows an increase in values of 1.78 times than control. Thus it can be concluded that the overall acceptability of fortified yogurt is better than the control in terms of sensory analysis, syneresis as well as functional properties.

Keywords: fortified yogurt, biocolour, purple cabbage, anthocyanin, functional foods.

I. Introduction

Functional foods can be obtained in whole or fortified, which provide health benefits by enriching essential nutrients (e.g., vitamins and minerals). Various functional foods have the potential to perform role in disease prevention. But, only a small number of these have had substantive clinical documentation of their health benefits [1]. Dairy products or the milk derived products can be considered as functional foods as they are health

beneficial for humans [2]. Functional dairy foods, the probiotics already have positive health image by their traditional use for centuries [3]. The market of functional food increased rapidly [2].

In plant foods, it is found that fruits and vegetables are rich in bio-active compound viz. phenolic compound, carotenoids, flavonoids, anthocyanin, vitamins, higher content of antioxidant and antimicrobial compounds. In fact the waste peels, seeds can be used as a source of nutraceutical. Thus higher intake of fruits and vegetables offers an effective tool for preventing serious disease like cancer, cardiovascular disease [4].

Colour is one of the most important characteristic of food. Main objective for the addition of colour in food is to make the food more appealing and recognizable. Everyone is sensitive to the colour of the food as it can stimulate or suppress one's appetite [5]. Natural colorants have received increased acceptability, mainly because of the apparent lack of toxicity. Natural colours are sometimes rich sources of antioxidants [6].

Purple cabbage (*Brassica oleracea* var. *capitata* F. *rubra*) is a vegetable with good antioxidant capacity. The leaves of this cabbage contain biologically potent anthocyanins [7]. It has anticancerous [8], anti-inflammatory [9] antibacterial [10], anti-diabetic properties [11] as well as it possesses antioxidant, and antihyperglycemic [12]. It is also very rich in minerals, vitamins, anthocyanins, polyphenols, and glucosinolates. It was observed that it contains about 196.5 mg of polyphenols per 100g of raw purple cabbage. It is more unique among the cruciferous vegetables as it provides a good quantity of anthocyanins, which acts as antioxidant as well as anti-inflammatory nutrients. The antioxidant richness of cabbage is partly responsible for its cancer prevention benefits [13] [14].

Anthocyanins are water soluble pigment and they are also known as flavonoids. They provides the bright and attractive orange, red, purple, and blue colours of most fruits, vegetables, flowers and some

Study of Combined Action of Papaya Leaf Extract, Pomegranate and Lemon – A Review

Dr.Bandyopadhyay¹ | Prithweejit Bhattacharyya² | Lopamudra Banerjee² | Pallabi Pal² | Shatarghya Das Mazumder²

¹H.O.D and Associate Professor, Department of Food technology, Guru Nanak Institute of Technology, Kolkata, India.

²UG Students, Department of Food technology, Guru Nanak Institute of Technology, Kolkata, India.

To Cite this Article

Dr.Bandyopadhyay, Prithweejit Bhattacharyya, Lopamudra Banerjee, Pallabi Pal and Shatarghya Das Mazumder, "Study of Combined Action of Papaya Leaf Extract, Pomegranate and Lemon – A Review", *International Journal for Modern Trends in Science and Technology*, Vol. 06, Issue 05, May 2020, pp.: 67-71; <https://doi.org/10.46501/IJMTST060511>

Article Info

Received on 24-March-2020, Revised on 24-April-2020, Accepted on 28-April-2020, Published on 02-May-2020.

ABSTRACT

The main motive of this review study is to focus on the essential components present in papaya leaves, pomegranate juice and sweet lime which is mausambi juice and a result of their combined effect on human health with enhanced beneficial components such as saponins, polyphenols etc. The leaves of papaya are generally considered as waste but their extracts have various health benefits. It contains calcium, magnesium, iron, and vitamins A, C, E, K and B. It is said that Saponin which is a component present in papaya leaves have anticancer capacity as well as reduces excessive oxidative stress. Here, papaya leaf extract can be added to pomegranate lemonade where both these fruits have predominant nutritional value. Pomegranate boosts immunity and aids in digestion. It also has anti-inflammatory properties. It is loaded with Vitamin-C and is also antioxidant-rich. Lemon on the other hand also contains a lot of minerals and vitamins, it also reduces oxidative stress. As a result it is an approach of a health drink having medicinal values. The positive effect of all the above extracts when combined can produce immense health benefits in the human body and can act as suitable nutraceutical with chances of less side effects and allergic symptoms.

KEYWORDS: Anti-inflammatory, antioxidant, immunity, nutraceutical, saponin.

Copyright © 2014-2020 International Journal for Modern Trends in Science and Technology

DOI: <https://doi.org/10.46501/IJMTST060511>

I. INTRODUCTION

The main aim of our study is to use *Carica papaya* leaf extract as a base to utilize its medicinal value and fortify it with commercially available pomegranate lemonade. papaya leaf extract has anti-cancer properties, added to that it also has a lot of health benefits such as- it may aid in digestion as it contains fiber, it has antioxidant properties, it may lower blood sugar level and may increase blood platelets count in dengue affected patients. Therefore, being cost effective, papaya leaf extract may solve a lot of health problems.

Carica papaya is usually cultivated in tropical regions, it's commonly called as papaya in this part of the world and it's from the family "Caricaceae". Papaya grows best in a well-drained, well aerated and rich organic matter soil, pH 5.5 – 6.7 (Priyanka et al. [1]), while water logging of soils often results in the death of trees within 3-4 days (Storey [2]). The leaves also contain active components such as papain, chymopapain, cysteine, ascorbic acid, flavonoids, cyanogenic glucosides that increases the total antioxidant power in blood and reduce lipid peroxidation level. Amount of total

Fortifications in Ice-cream with Enhanced Functional Properties: A Review

Dolanchapa Sikdar¹ | Rosalin Nath¹ | Kakali Bandyopadhyay¹ | Ishon Mollick¹

¹Department of Food Technology, Guru Nanak Institute of Technology, Kolkata, India

To Cite this Article

Dolanchapa Sikdar, Rosalin Nath, Kakali Bandyopadhyay and Ishon Mollick, "Fortifications in Ice-cream with Enhanced Functional Properties: A Review", *International Journal for Modern Trends in Science and Technology*, Vol. 06, Issue 05, May 2020, pp.: 131-138; <https://doi.org/10.46501/IJMTST060523>

Article Info

Received on 02-April-2020, Revised on 28-April-2020, Accepted on 05-May-2020, Published on 17-May-2020.

ABSTRACT

Enrichment of food by the addition of essential micro-nutrients like trace elements and vitamins to food is known as fortification. It can be carried out under initiatives such as Public Health Policy which aims in reducing the number of people with dietary deficiencies within a group of people or population. Ice-creams are well known dessert, and very much in demand among the children. So it is very important to keep its sensory factors in mind and make nutritive additions in it. To meet the demands of the consumers various natural ingredients are used to fortify ice-creams. Works have been done on combining ice-cream with biologically active molecules and produce functionally potential products. There are reports of making natural anti-oxidant ice-cream, experimenting it with cocoa powder, hazelnut, green-tea and ginger extract. Ice-cream fortification with fish-protein powder has turned to be effective in enhancing food value, but studies on its stability during storage and consumer acceptance are highly recommended. Researchers worked on incorporating dietary fibers in ice-cream like wheat, oats, bamboo, apple and orange. Its influence was checked on the basis of rheological, textural, sensory properties and probiotic culture survival of the ice-cream, to enhance the structural characteristics and cultural viability. Addition of Inulin in ice-creams as fat-replacers increased the viscosity and also made it resistant to melting. Hydrocolloids bind water and oil and adding them in ice-creams enhanced the micro-viscosity. Ice-creams were fortified with vitamins and minerals with successful attempts for example Vitamin D3, calcium and zinc fortifications. This study aims to review the various fortifications of ice-cream till date, using different nutritional components, hence fulfilling the nutritional needs of human-beings.

KEYWORDS: Fortification, Vitamins, Dietary deficiency, Biologically active molecules, Sensory properties, Micro-viscosity, Dietary fibers.

Copyright © 2014-2020 International Journal for Modern Trends in Science and Technology

DOI: <https://doi.org/10.46501/IJMTST060523>

I. INTRODUCTION

When any food product is developed, the first few things that must be kept in mind is fulfilling the consumers' expectancy and checking their acceptance of the product. It is a game of combining sensory properties of food with

nutritional factors. Any kind of deficiency in the daily dietary routine causes the development of technologically advanced food (enriched in essential micronutrients). Fortification may be defined as the enrichment of food by the addition of essential nutritive. The fortification of foods is often



ISSN (E): 2277- 7695
ISSN (P): 2349-8242
NAAS Rating: 5.03
TPI 2019; 8(12): 416-420
© 2019 TPI
www.thepharmajournal.com
Received: 19-10-2019
Accepted: 21-11-2019

Shairee Ganguly

Assistant Professor, Department of Food Technology, Guru Nanak Institute of Technology, Nilgunj Road, Sodpur, Panihati, Kolkata, West Bengal, India

Kakali Bandyopadhyay

Associate Professor and Head, Department of Food Technology, Guru Nanak Institute of Technology, Nilgunj Road, Sodpur, Panihati, Kolkata, West Bengal, India

Purba Dutta

B. Tech. Students, Department of Food Technology, Guru Nanak Institute of Technology, Nilgunj Road, Sodpur, Panihati, Kolkata, West Bengal, India

Anirzeet Pramanik

B. Tech. Students, Department of Food Technology, Guru Nanak Institute of Technology, Nilgunj Road, Sodpur, Panihati, Kolkata, West Bengal, India

Corresponding Author:

Shairee Ganguly

Assistant Professor, Department of Food Technology, Guru Nanak Institute of Technology, Nilgunj Road, Sodpur, Panihati, Kolkata, West Bengal, India

A review on microbial-pigment: A good source of biocolour

Shairee Ganguly, Kakali Bandyopadhyay, Purba Dutta and Anirzeet Pramanik

Abstract

In today's world there is a growing demand for healthy food products has lead to the development of "Nutraceutical Food Products". Development of natural food colours or bio-colours is a growing research area to shift the synthetic food colours. Bio-colours have many health benefits properties such as anti oxidant activity, food products stabilization, anti cancer property, etc. The main challenge faced by the industry is the production of natural food pigments in bulk to replace the synthetic colours and also this extraction process from plant origin is time consuming, expensive and the product yield is low. The sources are non-renewable. In this regard, the micro organisms are less expensive and renewable alternative source. In this review paper the competitive study of different microbial pigment is done as well as use of microbial pigment as an alternative of synthetic food pigment and the nutraceuticals properties of those pigments are discussed. This paper also highlights the challenges faced and advanced technologies used in this developing the pigment extracted from microbial sources.

Keywords: Bio-colour, micro-organism, nutraceutical, food pigments

1. Introduction

Colours are a main property of sensory evaluation of any food materials. Since ancient times, human have added food colours to the food products to make the food look more presentable, to make up for the loss of colours during processing and to improve the quality. Customers have since long associated different characteristics of food with colour such as freshness, taste etc. Hence colours have for long served as the sensory evaluating tool. Due to the ease of production in bulk and the cheapness of production, artificial food colours are being used in large scale. However the knowledge about the toxic effect of these artificial food colours among the population is coming to the fore front. Thus a market shift towards development and production of Biocolours has been observed. This shift is more so catalyzed by the recent development of Nutraceutical Food Products as the bio colours are known to contain many beneficial properties and also natural colours are eco friendly. Thus if they replace artificial colour, the production of harmful chemicals polluting the environment will also stop. Now biocolours are generally extracted from plant and vegetable sources which are non renewable sources. However it has been discovered that pigments extracted from micro organisms can be an alternate means of production of natural food colours. Hence a Seemingly vast number of micro organisms found in nature, ease of cultivation of these micro organisms and their medicinal properties adds to the advantage in extraction and production of natural colours from them. Bacteria, Fungi, Algae etc. all produces pigments which can be used for production of bio colours. Among the molecules produced by microorganisms are carotenoids, melanins, flavins, phenazines, quinines (Joshi *et al.*, 2003; Pankaj *et al.*, 2016) ^[6, 13].

2. Microbial pigments as food colours

In this earth, uncountable numbers of types of Micro organism are found. Many microbes such as micro algae, bacteria, fungi etc all are known to produce a variety of pigments. As colour extraction from microbial sources have some benefits like, cheaper production, easier extraction, higher yields through strain improvement, less lack of raw materials and no seasonal variations, microbial pigment has become a major source of bio colours (Kamla *et al.*, 2012; Dufosse *et al.*, 2009) ^[7, 3]. The health benefits of the microbial pigment such as anti oxidant, non carcinogenic, anti microbial etc adds to the use of microbes. *Flavobacterium sp.* (yellow pigment: zeaxanthin), *Agrobacterium aurantiacum* (pink- red pigment: astaxanthin), *Micrococcus sp.* (different colored pigments, carotenoids), *Serratia marcescens* (red pigment),

A REVIEW ON MUSHROOM: A CANCER ANTIDOTE

¹Rosalin Nath, ²Kakali Bandyopadhyay, ³Dolanchapa Sikdar and ⁴Arnab Saha

¹Assistant Professor, ²Associate Professor, ³Assistant Professor, ⁴Student
Department of Food Technology
Guru Nanak Institute of Technology, Panihati, Kolkata, India.

ABSTRACT: Cancer is the leading cause of death worldwide. The current anti-cancer drugs have numerous side effects. In a search for less toxic and effective treatment of cancer some prized mushrooms have been found to have validated anti-cancer properties. Numerous attempts have been made to assess their benefits for commercial purposes and cancer therapy. Certain bioactive molecules including polysaccharides, proteins, glycosides, alkaloids, volatile oils, tocopherols, phenolics, flavonoids, carotenoids, ascorbic acid enzymes, and organic acids which are also anti-tumour agents have been identified from various mushrooms. The anti-cancer potential of mushrooms lies in lentinan, krestin, hispolon, lectin, calcaelin, illudin S, psilocybin, Hericium polysaccharide A & B (HPA & HPB), ganoderic acid, schizophyllan, laccase, etc. Psilocybin mushrooms are used in folk medicine though it's not considered edible. Over 50 mushroom species have been found to yield immunoceuticals with anti-cancer properties for animals including six for human cancers. PSP (Polysaccharide-peptide) over the last 5 year tests have significantly improved quality of life with substantial pain relief besides enhancing immune status in most patients with cancers of stomach, oesophagus, lung, ovary, and cervix. PSK (Polysaccharid-K) and PSP boosted immune cell production, ameliorated chemotherapy symptoms, and enhanced tumours infiltration by dendritic and cytotoxic T-cells. These chemicals have extremely low side-effects, improve the quality of life, and highly compatible with chemo therapy and hence, well suited for cancer management regimens. So here in this review paper, anti-cancer drugs produced by edible & wild both mushrooms & their activity to prevent & cure cancer has been discussed briefly.

Keywords: Anti-cancer Drugs, Folk medicine, Immunoceuticals, cytotoxic T-cells, bioactive molecules.



ISSN (E): 2277- 7695
 ISSN (P): 2349-8242
 NAAS Rating: 5.03
 TPI 2019; 8(11): 207-211
 © 2019 TPI
 www.thepharmajournal.com
 Received: 14-09-2019
 Accepted: 18-10-2019

Anindya Dhar
 B.Tech 2nd year Student,
 Department of Food Technology,
 Guru Nanak Institute of
 Technology, 157/F, Nilgunj
 Road, Sodepur, Panihati,
 Kolkata, West Bengal, India

Poulami Chakraborty
 B.Tech 2nd year Student,
 Department of Food Technology,
 Guru Nanak Institute of
 Technology, 157/F, Nilgunj
 Road, Sodepur, Panihati,
 Kolkata, West Bengal, India

Shalini Chanda
 B.Tech 2nd year Student,
 Department of Food Technology,
 Guru Nanak Institute of
 Technology, 157/F, Nilgunj
 Road, Sodepur, Panihati,
 Kolkata, West Bengal, India

Dr. Anju Paul
 Assistant Professor, Department
 of Food Technology, Guru
 Nanak Institute of Technology,
 157/F, Nilgunj Road, Sodepur,
 Panihati, Kolkata, West Bengal,
 India

Corresponding Author:
Anindya Dhar
 B.Tech 2nd year Student,
 Department of Food Technology,
 Guru Nanak Institute of
 Technology, 157/F, Nilgunj
 Road, Sodepur, Panihati,
 Kolkata, West Bengal, India

A review on extraction of glucose from rice

Anindya Dhar, Poulami Chakraborty, Shalini Chanda and Dr. Anju Paul

Abstract

Carbohydrates are one of the most essential macro-nutrient required for the maintenance of life. Glucose and Fructose act as building blocks of energy. The Human Body consumes carbohydrates in the form of glucose. Glucose is converted to glycogen, a polysaccharide and it acts as a readily available source of energy. Approximately 130 g of glucose is required to nourish the brain cells throughout the day. Glucose is widely used in food manufacturing industries which include beverage, ice cream, alcohol, confectionary and other fermentation plants.

Studies and researches have shown that extraction of glucose from rice is a cost effective method. Rice is widely consumed and important cereal crop. The three variants of Indian rice HMT, Swarna and basmati have high starch contents. In a nutshell the paper is focused to concentrate about the different methods of glucose production from rice.

The major researches included in the paper are:

1. Glucose obtained from Rice bran by Ultrasound-Assisted enzymatic hydrolysis
2. Isolation of starch from Rice (*Oryza Sativa* L.) by alkali extraction method
3. Efficient Recovery of Glucose via enzymatic saccharification of rice straw with soft carbohydrates
4. Glucose production from rice husk by solid state fermentation method.

Keywords: Rice starch, enzymatic hydrolysis, rice amylase, Solid state fermentation, Saccharification

Introduction

Rice starch isolation is a bit different from other sources of extraction because of its unique protein composition. The isolation process mainly consists of the separation of other components like protein, fiber, and lipid. The key factor that one has to keep in mind is that the mechanical damage or amyolytic destruction of the starch granules must be avoided.




Fig: Steps of Rice Starch extraction

A Relative Study on Utilisation of Fenugreek Seeds for Enhancement of the Antioxidant Activities in Various Baked Products

Kakali Bandyopadhyay^{#1}, Shairee Ganguly^{#2}, Chaitali Chakraborty^{*3}, and Rupsa Roychowdhury^{#4}

^{#1}Associate Professor and HOD, ^{#2} Assistant Professor, ^{#4} Student

[#]Department of Food Technology, Guru Nanak Institute of Technology, 157/F, Nilgunj Road, Sodepur, Panihati, Kolkata 70011, West Bengal, India.

^{*3}Associate Professor, Department of Dairy Chemistry, Faculty of Dairy Technology, WBUAFS, Mohanpur, Nadia, West Bengal, India

Abstract

Fenugreek (*Trigonella foenum-graecum* L.) is a widely distributed well-known annual plant and has possessed obvious hypoglycaemic and hypercholesterolemia characteristics. In the present study fenugreek seeds and its extracts are utilized for fortification of low nutritional value baked items so as to enhance its nutraceutical properties. It was found that hot extraction and roasted & grinded fenugreek seed extracts had more polyphenol content than cold extraction and raw & grinded fenugreek seed. So it can be concluded that the amount of soluble polyphenols present in fenugreek seeds are increased after heat treatments. Result shows that, the maximum amount of polyphenols is present in three different types of baked items (bread: 0.304 ± 0.002 , cookies: 0.291 ± 0.003 , muffin: 0.345 ± 0.001 gm GAE/100gm sample) when fortified with hot extraction of fenugreek seeds.

Therefore, for fortification of heat treated baked food items like bread, cookies and muffins, fenugreek seed and its extracts are good sources of polyphenols.

Key words: Fenugreek, polyphenol, heat stable, baked food, nutraceutical

I. INTRODUCTION

Fenugreek (*Trigonella foenum-graecum* L.) is a legume and it has been used as a spice throughout the world to enhance the sensory quality of foods. Fenugreek is well-known for its medicinal properties such as anti-diabetic, anti-carcinogenic, hypocholesterolemic, antioxidant, and immunological activities.

Studies have shown fenugreek seeds to be also a rich source of antioxidants. Polyphenols in fenugreek seeds include apigenin and a number of kaempferol and quercetin glycosides [1] as well as flavonoids; vitexin, tricetin, naringenin quercetin and tricetin-7-O- β -D-glucopyranoside (Shang *et al.*, 1998). The extracts of endosperm husk, and fenugreek seed at about 200mg

concentration exhibited antioxidant activity 72%, 64%, and 56% respectively by free-radical scavenging method [2]. The major bioactive compounds in fenugreek seeds are believed to be polyphenol compounds, such as rhaponticin and isovitexin [3]. Dixit *et al.* (2005) [4] found that fresh leaves of fenugreek contain ascorbic acid of about 220.97 mg per 100 g of leaves and β -carotene is present about 19 mg/100 g. On the other side, it was reported that 84.94% and 83.79% ascorbic acid were reduced in sun and oven-dried fenugreek leaves respectively.

Health benefits of fenugreek seeds include:

- Diabetes Mellitus prevention: Fenugreek powder treatment in patients suffering from mild Non-insulin dependent diabetes mellitus produced marked reduction in blood sugar and serum triglycerides and total cholesterol [5].
- Cancer prevention: The effect of fenugreek seeds observed in induced breast cancer [6]. Further, the ethanolic extract of fenugreek showed antineoplastic effect on the growth of breast cancer cells by reducing cell viability, inducing early apoptotic changes, declining the mitochondrial membrane potential and degrading cellular DNA into fragments [7].
- Antioxidant activity: It has been documented in various studies that fenugreek bears potential of a powerful antioxidant in which the presences of flavonoids and polyphenols have been found to be responsible for the same [4]. The exposure of polyphenol rich extract of fenugreek seeds which showed protective effects against hydrogen peroxide induced oxidation by protecting the erythrocytes from haemolysis and lipid peroxidation in a dose dependent manner [8].

This study was carried out with the objective to enhance the antioxidant properties of various baked products made from soft wheat flour on fortification



ISSN (E): 2277- 7695

ISSN (P): 2349-8242

NAAS Rating: 5.03

TPI 2019; 8(5): 397-398

© 2019 TPI

www.thepharmajournal.com

Received: 10-03-2019

Accepted: 11-04-2019

Dr. Anju Paul

Assistant Professor, Department of Food Technology, Guru Nanak Institute of Technology, 157/F, Nilgunj Road, Sodepur, Panihati, Kolkata, West Bengal, India

Debarati Bhattacharyya

B.Tech Final Year Student, Department of Food Technology, Guru Nanak Institute of Technology, 157/F, Nilgunj Road, Sodepur, Panihati, Kolkata, West Bengal, India

Analysis and study on the antioxidant of citrus fruits

Dr. Anju Paul and Debarati Bhattacharyya

Abstract

The term "antioxidant" is mainly used for two different groups of substances: industrial chemicals which are added to products to prevent oxidation, and natural chemicals found in foods and body tissue which are said to have beneficial health effects. Citrus fruits are the source of the second category. They are fruits of the genus *Citrus* having thick rind and juicy pulp. *Citrus Fruits* are well-known for its nutrition and health-promotion values. In recent years, the antioxidant activity of *Citrus* fruits and their roles in the prevention and treatment of various human chronic and degenerative diseases have attracted more and more attention. *Citrus* fruits are suggested to be a good source of dietary antioxidants. To have a better understanding of the mechanism underlying the antioxidant activity of *Citrus* fruits, we have done these experiments of analysing acidity, moisture content & antioxidant content of 3 citrus fruits (will be discussed below).

Keywords: Citrus fruits, antioxidants, acidity of fruits, health benefits of fruits

Introduction

Citrus is a genus of trees with fleshy, juicy fruits, widely used edible plants belonging to *Citrus* and related genera of the family *Rutaceae* (orange family). Included are the tangerine, citrange, tangelo, orange, pomelo, grape fruit, lemon, lime, citron, and kumquat. Introduced throughout Europe during the Crusades, they were brought by Portuguese and Spanish explorers to the West Indies, hence they were introduced into North and South America. Commercially they are now the most important group of tropical and subtropical fruits in the world. The fruits are rich in vitamin C (ascorbic acid), various fruit acids (especially citric acid), and fruit sugar. The rind, which contains numerous oil glands, and the fragrant blossoms of some species are also a source of essential oils used for perfumes and similar products. Citrus fruits can be damaged by freezing temperatures, pests (scale insects, rust mites), and various bacterial, viral, and fungal diseases (e.g., citrus canker, greening, tristeza, and melanose).

How can antioxidants benefit our health?

Antioxidants are substances that can prevent or slow damage to cells caused by free radicals, unstable molecules that the body produces as a reaction to environmental and other pressures.

They are sometimes called "free-radical scavengers."

Free radicals are waste substances produced by cells as the body processes food and reacts to the environment. If the body cannot process and remove free radicals efficiently, oxidative stress can result. This can harm cells and body function. Free radicals are also known as reactive oxygen species (ROS).

Antioxidants act as radical scavenger, hydrogen donor, electron donor, peroxide decomposer, singlet oxygen quencher, enzyme inhibitor, synergist, and metal-chelating agents. Antioxidant supplements may help reduce vision loss due to age-related macular degeneration in older people.

Foods that are particularly high in antioxidants are often referred to as a "superfood" or "functional food."

Health benefits of Citrus fruits are associated with its high amounts of photochemical and bioactive compounds such as flavonoids, limonoids, phenols, carotenoids, minerals and vitamins. The plant is used in traditional medicine as an antiseptic, antiviral, antifungal, astringent, diuretic, mosquito bite repellent, for the treatment of stomach ailments.

Materials & methods

Three types of Citrus fruits, Lemon (*Citrus lemon*), Sweet lime or Mosambi (*Citrus limetta*) Orange (*Citrus sinensis*) were taken.

Correspondence

Dr. Anju Paul

Assistant Professor, Department of Food Technology, Guru Nanak Institute of Technology, 157/F, Nilgunj Road, Sodepur, Panihati, Kolkata, West Bengal, India



ISSN (E): 2277- 7695

ISSN (P): 2349-8242

NAAS Rating: 5.03

TPI 2018; 7(9): 337-342

© 2018 TPI

www.thepharmajournal.com

Received: 09-07-2018

Accepted: 10-08-2018

Anju Paul

Department of Food Technology,
Guru Nanak Institute of
Technology, Panihati, Kolkata-
700 114, West Bengal, India

Mainak Debnath

Department of Applied Science &
Humanities, Guru Nanak
Institute of Technology,
Panihati, Kolkata-700 114,
West Bengal, India

Mainak Debnath

Department of Applied Science &
Humanities, Guru Nanak
Institute of Technology,
Panihati, Kolkata-700 114,
West Bengal, India

Amit Kumar Barman

Faculty of Dairy Technology,
WBUAFS, Mohanpur campus,
West Bengal, Nadia

A veteran vitamin in versatile role: Vit K

Anju Paul, Sreyasree Basu, Mainak Debnath and Amit Kumar Barman

Abstract

Vitamin K is precisely known for its role in blood clotting. The original term vitamin “K” comes from the K in the Germanic word Koagulation meaning the ability to clot blood or prevent hemorrhage. In recent days, new roles for vitamin K have been emerged in the multifarious fields like, brain health, bone, cardiovascular, hormonal and reproductive health etc. vitamin K deficiency is much more likely to occur in infants but in adults person it is also seen. Symptoms include bruising, hematomas oozing of blood at surgical or puncture sites, stomach pains; risk of massive uncontrolled bleeding; cartilage calcification; and severe malformation of developing bone or deposition of insoluble calcium salts in the walls of arteries. In infants, it can cause some birth defects such as underdeveloped face, nose, bones, and fingers. Vitamin K is changed to its active form in the liver by the enzyme Vitamin K epoxide reductase. Activated vitamin K is then used to gamma carboxylate (and thus activate) certain enzymes involved in coagulation: Factors II, VII, IX, X, and protein C and protein S. Inability to activate the clotting cascade via these factors leads to the bleeding symptoms.

Keywords: Vitamin K, coagulation, calcification, phylloquinone, menaquinones, menadiione

Introduction

The discovery of vitamin K can be traced back to the research of Carl Peter Henrik Dam at the Biochemical Institute of the University of Copenhagen from 1928 to 1930. In his work on cholesterol metabolism, the Danish biochemist observed a spontaneous tendency to hemorrhage in chicks fed for longer than 2 to 3 weeks on cholesterol- and fat-free chicken feed. This coagulation disorder was combined with lowered prothrombin content (Prothrombin D factor II) of the blood [1-3]. At that time, as none of the hitherto known vitamins (e.g. vitamins A, C and D) were capable of preventing the coagulation disorder, Dam postulated a new, fat-soluble vitamin, which regulates coagulation. The latter was apparently present in green vegetables and liver, as supplementary feeding with these nutrients resulted in normal blood coagulation in the animals. Moreover, Dam successfully treated the chickens' hemorrhages with an ether extract obtained from lucerne (alfalfa). Dam Called the antihemorrhagic vitamin “vitamin K” (after “Koagulation:” coagulation) [4]. In the 1930s, several working groups researched the isolation and identification of vitamin K. At this time, a US American research group working with the biochemist Edward Albert Doisy succeeded in isolating the antihemorrhagic vitamin K and elucidating its chemical naphthoquinone ring structure. In 1943, the 2 researchers, Dam und Doisy, were jointly awarded the Nobel Prize for medicine for the discovery and elucidation of the chemical structure of vitamin K [5, 6].

Vitamin K is a name given to a group of fat-soluble vitamins. They are considered essential cofactors in humans for the production of several proteins that are involved in coagulation homeostasis and calcium homeostasis.

Chemically, the vitamin K family comprises 2-methyl-1,4-naphthoquinone (3-) derivatives. Vitamin K includes two natural vitamers: vitamin K₁ and vitamin K₂. Vitamin K₂, in turn, consists of a number of related chemical subtypes, with differing lengths of carbon side chains made of isoprenoid groups of atoms.

Vitamin K₁, also known as phylloquinone, is made by plants, and is found in highest amounts in green leafy vegetables because it is directly involved in photosynthesis. It may be thought of as the plant form of vitamin K. It is active as a vitamin in animals and performs the classic functions of vitamin K, including its activity in the production of blood-clotting proteins. Animals may also convert it to vitamin K₂.

Bacteria in the gut flora can also convert K₁ into vitamin K₂ (Menaquinone). In addition, bacteria typically lengthen the isoprenoid side chain of vitamin K₂ to produce a range of vitamin K₂ forms, most notably the MK-7 to MK-11 homologues of vitamin K₂.

Correspondence

Anju Paul

Department of Food Technology,
Guru Nanak Institute of
Technology, Panihati, Kolkata-
700 114, West Bengal, India



ISSN (E): 2277- 7695

ISSN (P): 2349-8242

NAAS Rating: 5.03

TPI 2019; 8(1): 42-47

© 2019 TPI

www.thepharmajournal.com

Received: 26-11-2018

Accepted: 30-12-2018

Riya Dasgupta

Quality Assurance Manager,
Agrion Overseas Private
Limited, Sangli, Maharashtra,
India

Chaitali Chakraborty

Associate Professor, Department
of Dairy Chemistry, Faculty of
Dairy Technology, WBUAFS,
Mohanpur, Nadia, West Bengal,
India

Kakali Bandyopadhyay

Head & Associate Professor,
Department of Food Technology,
Guru Nanak Institute of
Technology, Panihati, Sodepur,
Kolkata, West Bengal, India

Shairee Ganguly

Assistant Professor, Department
of Food Technology, Guru
Nanak Institute of Technology,
Panihati, Sodepur, Kolkata,
West Bengal, India

Development and characterization of biocolor (*Brassica oleracea var. capitata F. rubra*) fortified lime squash

Riya Dasgupta, Chaitali Chakraborty, Kakali Bandyopadhyay and Shairee Ganguly

Abstract

The major pigment of purple cabbage is anthocyanin, responsible for many of the attractive colors, from scarlet to blue. It represents about 196.5 mg of polyphenols per 100g of raw purple cabbage that can be determined spectrophotometrically. Purple cabbage also contains high amount of dietary fiber, carbohydrates and related bioactive compounds. It is thus significant and even essential to find applications in different food products (e.g. Lime Squash). Squash is a non-alcoholic concentrated syrup used in beverage making. It is usually fruit-flavored, made from fruit juice, water, and sugar or a sugar substitute. In this study color extracted from purple cabbage was used for the development of lime squash, packed in glass bottles and sealed hermetically and non-hermetically using wax. The polyphenol of all samples were evaluated spectrophotometrically and expressed in mg Gallic acid equivalent/gm of sample. The light absorbance and color stability of the sample were studied using UV-Vis double-beam spectrophotometer. In this study, the peak of anthocyanin is processed in between 500-600 nm and the maximum peak height of anthocyanin is found at 560nm and color stability is determined within the wavelength range between 550-560nm. The best result in all aspects was found for the lime squash sealed hermetically.

Keywords: purple cabbage, anthocyanin, lime squash, gallic acid equivalent, color stability

1. Introduction

Color is a measure of quality and nutrient content of foods. The objective of adding color to foods is to make them appealing, augment the loss of color during processing, to improve the quality and also to influence the consumer to buy a product. At present, the demand for natural dyes is increasing worldwide due to the increased awareness on therapeutic and medicinal properties and their benefits among public and also because of the recognized profound toxicity of synthetic colors. Natural dyes are those derived from naturally occurring sources such as plants, insects, animals and minerals. Among all the natural dyes, plant-based pigments have medicinal values so are mostly preferred (Chaitanya, 2014). The major pigments present abundantly in purple cabbage are anthocyanin which can be used as a source of natural color. The term anthocyanin was derived from the Greek *anthos*, a flower, and *kyanos*, dark blue. Anthocyanins are the most important group of pigments, after chlorophyll that is visible to the human eye (Harborne *et al.*, 1988). Anthocyanins are responsible for many of the attractive colors, from scarlet to blue, of flowers, fruits, leaves, and storage organs (Harborne *et al.*, 1988-1993). Being part of flavonoids, anthocyanins are the greatest natural pigments giving blue color in plants and showing antioxidant potential.

Purple cabbage belonging to *Brassicaceae* family is a vegetable with high total antioxidant capacity being also very rich in minerals, vitamins, polyphenols, anthocyanins and glucosinolates. It represents about 196.5 mg of polyphenols per 100g of raw purple cabbage. Purple cabbage also contains high amount of dietary fiber, carbohydrates and related bioactive compounds. Purple cabbage, is more unique among the cruciferous vegetables in providing a big quantity of anthocyanins, which qualify not only as antioxidant nutrients, but also as anti-inflammatory nutrients. The antioxidant richness of cabbage is partly responsible for its cancer prevention benefits (Draghici *et al.*, 2013) ^[5].

2. Materials and Methods

2.1 Materials

2.1.1 Raw Materials Required: Raw materials for the preparation of this squash such as purple cabbage, sugar, limes were brought from local market adjacent to the institution.

Correspondence

Riya Dasgupta

Quality Assurance Manager,
Agrion Overseas Private
Limited, Sangli, Maharashtra,
India

Studies on extraction of polyphenols from food wastes and its utilization for fortification of polyphenols

■ **Kakali Bandyopadhyay, Shairee Ganguly, Chaitali Chakraborty and Anusree Modak**

Received : 01.12.2017; Revised : 06.03.2018; Accepted : 18.03.2018

See end of the Paper for authors' affiliation

Correspondence to :

Kakali Bandyopadhyay
Department of Food
Technology, Gurunanak
Institute of Technology,
Panihati, Sodepur, **Kolkata**
(W.B.) India
Email : head_ft.gnit@jsgroup.org

■ **ABSTRACT** : Food processing is one of the most important industries over the world, however, byproducts of such industries, mainly organic material must be handled in appropriate manner to avoid any environmental violence. But these by products as well as the waste materials contain some good nutraceutical properties such as presence of polyphenols which can increased the antioxidant properties of the food materials. In this study the polyphenol contents from the wastes of some food materials such as cucumber peels, potato peels, pea nut hulls, pomegranate seed wastes, date seeds, pumpkin seeds are determined in mg GAE/g sample by using different solvents viz., water, ethanol, methanol and acetone as extracting medium. It was observed that, among these samples potato peels, cucumber peels and pomegranate seeds exhibited maximum polyphenol content in water medium, such as 5.14, 6.05 and 4.70 mg GAE/g sample, respectively, whereas pumpkin seeds and date seeds showed maximum polyphenol content in acetone medium, like, 6.06 and 4.40 mg GAE/ g sample, respectively. As water is one of the main ingredients of most of the processed foods, potato peels and cucumber peels wastes can be utilized for fortifications of polyphenols in vegetable soups. It has been observed that the polyphenol content can be increased upto about 2.7 times in case of cucumber peel fortified soups. Therefore, it can be concluded that cucumber peel fortified soups showed the best result considering its functional quality in terms of polyphenol content as well as overall acceptability in terms sensory analysis.

■ **KEY WORDS** : Food wastes, Polyphenol, Nutraceutical, Fortified soups, Sensory analysis

■ **HOW TO CITE THIS PAPER** : Bandyopadhyay, Kakali, Ganguly, Shairee, Chakraborty, Chaitali, Modak, Anusree (2018). Studies on extraction of polyphenols from food wastes and its utilization for fortification of polyphenols. *Internat. J. Agric. Engg.*, **11**(1) : 216-219, DOI: 10.15740/HAS/IJAE/11.1/216-219.

BANANA PEEL WASTES: POTENTIAL SOURCE OF ANTIOXIDANT IN BANANA CHIPS

CHAITALI CHAKRABORTY¹, KAKALI BANDYOPADHYAY²,
BORNINI BANERJEE³ & SHUBHAM MUKHERJEE⁴

^{1,2}Assistant Professor, Department of Food Technology, Guru Nanak Institute of Technology, Panihati, Sodepur, Kolkata, India

^{3,4}Student, Department of Food Technology, Guru Nanak Institute of Technology, Panihati, Sodepur, Kolkata, India

ABSTRACT

Banana are one of the most popular fruit carries a number of beneficial pharmacological effects and comes with a set of variety and it is distributed all over the world. The major by-product of the banana processing industry is the peel causing environmental hazard. High dietary fibre, polyphenolic and related bioactive compounds content of banana peels make them promising for variety of applications in nutraceuticals and medicinal purposes.

In the present study the banana chips with and without peel were prepared by drying (using tray dryer at 60°C, 70°C, 80°C) and microwave processing (360, 540, 720, 900 watt). The results showed that moisture loss and shrinkage loss of banana chips increased with increase in time of drying and temperature and wattage of microwave. All types of chips were accepted by sensory analysis with highest gradation was obtained by microwave processed chips without frying. Polyphenol content (mg GAE/gm) of tray dried and microwave dried banana chips with and without peel were investigated which showed that microwave processed banana chips with peel contain highest amount of polyphenols compared to other varieties. Hence consumption of microwave processed banana chips with peel may be useful to combat free radical mediated diseases.

KEYWORDS: Banana Chips, Peel, Polyphenol, Microwave, Sensory Analysis & Tray Drier

Received: Jan 09, 2017; **Accepted:** Feb 11, 2017; **Published:** Feb 20, 2017; **Paper Id.:** IJASRAPR201711

INTRODUCTION

In ancient Indian literature the medicinal properties of banana have been documented and found to be effective in cure of many diseases [1]. As a staple fruit, it is available through-out the year which provides livelihood security to thousands of people [2]. In banana processing industry the major waste is the peel, accounting 30% of the fruit which constitute environmental hazard [3]. According to the criteria established by the National Cancer Standard Institute, banana peel extract is classified as non-toxic to normal human cells [4]; therefore, it can be safely utilized as a natural source of antioxidants for value addition. It grows in humid low land to upland tropical areas [5]. Being as a tropical plant, banana protects itself from the oxidative stress caused by strong sunshine and high temperature by producing large amounts of antioxidants [6].

The peel and pulp of fully ripe banana provides good antimicrobial and antibiotic properties [7]. Banana peel is also an underutilized source of phenolic compounds. It accounts 40% of the total weight of fresh banana and these are used as fertilizer or discarded in many countries [8]. It contains various antioxidant compounds such as galocatectin and dopamine and it is also the rich source of total phenolics and this in turn reflects their

Original Research Article

<https://doi.org/10.20546/ijcmas.2017.604.281>

Textural Analysis of Spongy Indian Milk Dessert (Rasogolla) Fortified with Potato Powder

Chaitali Chakraborty* and Kakali Bandyopadhyay

Department of Food Technology, Guru Nanak Institute of Technology, 157/F, Nilgunj Road,
Panihati, Sodepur, Kolkata-700114, West Bengal, India

*Corresponding author

ABSTRACT

Potato is probably the most popular food item in the Indian diet and India is one of the largest producers of potato. Potato is a very rich source of starch. It also contains phosphorus, calcium, iron and some Vitamins C and A. It is vastly consumed as a vegetable and is also used in various forms such as starch, flour, alcohol and dextrin. Besides being used as a daily food item in various vegetable preparations, potato today increasingly finds use in the form of chips or wafers as snacks food. In the present study, potato powder is produced from two varieties of potato like kufri chandramukhi and kufri jyoti. These potato powders are utilized as a binding agent to produce fortified rasogolla sample R1 (kufri jyoti fortified) and R2 (kufri chandramukhi fortified). Control sample (C) of rasogolla is prepared using chhana (a heat and acid coagulated milk protein mass and an Indian equivalent to cottage cheese) only and a market sample (M) is also taken into consideration during comparative analysis of both sensory and texture. It has been found that the highest overall acceptability of about 8.7 (based on 9-point hedonic scale) is observed for sample R2 compared with others. Whereas considering the textural profile like hardness, fracturability, springiness and cohesiveness the sample R2 provides comparable result with other control and market sample.

Keywords

Potato powder, Rasogolla, Texture analysis, Sensory analysis, Chhana

Article Info

Accepted:
20 March 2017
Available Online:
10 April 2017

Introduction

Today potatoes have become an integral part of much of the world's cuisine and are the world's fourth-largest food crop, following rice, wheat, and maize. China is now the world's largest potato-producing country, and nearly third of the world's potatoes are harvested in China and India. Potatoes have high nutritive value and can be consumed in various forms and one of these is potato powder [International Year of the Potato 2008 – The potato (2009)]. Processing of potato into flour is the most satisfactory method of creating a product that is not only

functionally adequate, but also remain for an extended period without damage. Potato flour can become a highly viable value added product due to its versatility in function as a thickener and color or flavor improver (Avula *et al.*, 2006; Raj *et al.*, 2008). Several products are prepared by incorporating potato flour with other flours and using the processes like baking, roasting, steaming, boiling and deep fat frying (Gahlawat and Sehgal, 1996).

It also imparts a distinctive, pleasing flavour and improves toasting qualities and can be

Original Research Article

<https://doi.org/10.20546/ijcmas.2017.604.263>

Development and Characterization of Biocolour (*Beta vulgaris*) Enriched Low Calorie Lassi (Yogurt Based Beverage)

Shairee Ganguly, Chaitali Chakraborty* and Kakali Bandyopadhyay

Department of Food Technology, Guru Nanak Institute of Technology, 157/F, Nilgunj Road, Panihati, Kolkata-700114, West Bengal, India

*Corresponding author

ABSTRACT

Lassi is a popular traditional yogurt based drink from the Indian Subcontinent. Lassi is a blend of yogurt, water, spices and sometimes fruits. To increase the physical acceptability of lassi, it has been fortified with betalain powder which provides reddish colour. Colour is an important parameter for sensory analysis and consumer preference. However using synthetic colour could be harmful for health. In today's progressive world a shift from synthetic to biocolour is observed. Betalain is a biocolour extracted from beet-root (*Beta vulgaris*). It also increases the polyphenol content of the product which will increase the anti-oxidant property of this beverage. In the present study, effort has been given to reduce the calories of lassi by using zero calorie sweetener, sucralose as it is 600 times sweeter than sucrose. To avoid the heat damage of biocolour, freeze dried powder is used as colorant. The polyphenol content of betalain powder is measured in gallic acid equivalents (GAE) by Folin-Ciocalteu assay using different solvents like methanol, ethanol, petroleum ether and water. Polyphenol content is increased from 755.32 gm/100gm for control lassi to 862.1 gm/100gm sample for fortified lassi. Proximate analysis of the lassi samples and the stability of betalain are also determined. Finally, the sensory evolution is also executed by using 9 point hedonic scale. Thus fortified lassi is prepared with enhanced functional properties as well as its overall acceptability.

Keywords

Lassi, Betalain, Polyphenol, Freeze-drying, 9 point hedonic scale, Proximate analysis.

Article Info

Accepted:
20 March 2017
Available Online:
10 April 2017

Introduction

India is diversified country, famous for its tradition, culture in each field because of agro-climatic zone, language, etc. India is popular for its traditional beverages and foods. These traditional foods are significantly known for their taste and texture, nutritional and therapeutic value. The combination of this traditional knowledge with scientific knowledge will increase the demand of some fermented foods to the health conscious consumers. For example, lassi is a

popular traditional yogurt based drink from the Indian Subcontinent. Lassi is a blend of yogurt, water, spices and sometimes fruits. In this present study, to increase the physical acceptability of lassi, it has been fortified with betalain (beet root color).

Again, effort has been given to reduce the calories of lassi by using non-nutritive sweetener, sucralose as it is 600 times sweeter than sucrose and has no calorie.

RESEARCH ARTICLE

Evaluation of rheological, physicochemical, and sensory properties of apple incorporated yoghurt

Chaitali Chakraborty, Suparna Mukherjee and Srijanee Biswas

Received: 05 May 2018 / Accepted: 19 December 2018 / Published online: 21 February 2019

© Indian Dairy Association (India) 2019

Abstract: Yoghurt, one of the fermented milk products, is a functional food which is popular and acceptable by consumers for several years. Yoghurt is a high nutritious food containing α -lactalbumin, β -lacto globulin, vitamin A, calcium and phosphorus. It is used to aid a variety of gastrointestinal disorders. Apples have higher shelf life (1-2 months in refrigerator) compared to many other fruits. Apples consist higher amount of fructose (about 12.6 gms) that acts as a natural sweetener which decreases the requirement of added sucrose to yoghurt. In this study a set-type fruit yoghurt was developed by addition of apple pulp. The physico-chemical, sensory and rheological properties of different types of yoghurt were determined. A control (C) was prepared to compare the quality characteristics with Apple Incorporated Yoghurt. The pH of the apple incorporated yoghurt lies between 3.89-4.01 whereas for control yoghurt it is between 4.14-4.34. The addition of apple pulp lead to decrease the pH, syneresis percentage, viscosity and an increase in total solid (TS), water holding capacity (WHC) of the yoghurt. AY3 revealed the highest () WHC (%) whereas the lowest value was observed in case of C. Flow index behavior for C, AY1, AY2 and AY3 were found as 0.57, 0.418, 0.402 and 0.395 respectively. So it can be concluded that

apple incorporated yoghurt serves as an excellent food worldwide and can be easily accepted by the consumers.

Keywords: Apple incorporated yoghurt, Physico-chemical analysis, Rheological analysis, Sensory analysis

Introduction

Yoghurt, which is a commonly fermented dairy product, is obtained from the fermentation of milk by lactic acid-producing bacteria such as *Streptococcus thermophilus* and *Lactobacillus delbrueckii spp. bulgaricus* (Fadela *et al.*, 2009). Yoghurt has a variety of probiotics and vitamins and is well received by consumers. Consumption of yoghurt is associated with some health benefits because of the lactic acid bacteria. As there is a lot of sugar and fat in it, it is not suitable for hyperpietics and diabetics. In recent years, consumers' demand for low-fat or nonfat dairy products has increased (Bitaraf *et al.*, 2012).

Yogurt is considered as healthy food due to its high digestibility and bioavailability of nutrients and also can be recommended to the people with lactose intolerance, gastrointestinal disorders such as inflammatory bowel disease and irritable bowel disease, and aids in immune function and weight control (Danone, 2013). Addition of flavors would enhance the consumer appeal while produce a variety of products. Flavors can either be added immediately before homogenization or after the homogenization. Yogurts are available in a vast array of flavors including fruit (apple, apricot, black cherry, black currant, blue berry, lemon, mandarin, raspberry, strawberry, peach), cereal, vegetables, chocolate, vanilla, caramel, ginger, etc (Weerathilake *et al.*, 2014; Chakraborty *et al.*, 2017).

Materials and methods

Fruit pulp preparation

Mature ripened apples were brought from local market of Sodepur, West Bengal and washed thoroughly. Then the apples were boiled at 90-95°C for around 5-10 minutes. After boiling, the skin, core and seeds of the apples were removed properly. Then

Chaitali Chakraborty (✉)
Department of Dairy Chemistry, Faculty of Dairy Technology,
WBUAFS, Mohanpur, Nadia, West Bengal, India
E-mail: ccty09@gmail.com

Suparna Mukherjee, Srijanee Biswas
Department of Food Technology,
Guru Nanak Institute of Technology, Panihati, Sodepur, Kolkata,
India,



ISSN (E): 2277- 7695
 ISSN (P): 2349-8242
 NAAS Rating 2017: 5.03
 TPI 2017; 6(10): 40-43
 © 2017 TPI
 www.thepharmajournal.com
 Received: 09-08-2017
 Accepted: 10-09-2017

Chaitali Chakraborty

HOD & Assistant Professor,
 Department of Food Technology,
 Gurunanak Institute of
 Technology, 157/F, Nilgunj
 Road, Panihati, Sodepur,
 Kolkata 700114, West Bengal,
 India

Kakali Bandyopadhyay

Associate Professor, Department
 of Food Technology, Gurunanak
 Institute of Technology, 157/F,
 Nilgunj Road, Panihati,
 Sodepur, Kolkata 700114, West
 Bengal, India.

Shairee Ganguly

Assistant Professor, Department
 of Food Technology, Gurunanak
 Institute of Technology, 157/F,
 Nilgunj Road, Panihati,
 Sodepur, Kolkata 700114, West
 Bengal, India.

Bornini Banerjee

Final year B. Tech Student
 Department of Food Technology,
 Gurunanak Institute of
 Technology, 157/F, Nilgunj
 Road, Panihati, Sodepur,
 Kolkata 700114, West Bengal,
 India.

Shubham Mukherjee

Final year B. Tech Student
 Department of Food Technology,
 Gurunanak Institute of
 Technology, 157/F, Nilgunj
 Road, Panihati, Sodepur,
 Kolkata 700114, West Bengal,
 India.

Correspondence**Chaitali Chakraborty**

HOD & Assistant Professor,
 Department of Food Technology,
 Gurunanak Institute of
 Technology, 157/F, Nilgunj
 Road, Panihati, Sodepur,
 Kolkata 700114, West Bengal,
 India

Potential of raw banana peel as a source of polyphenol in muffins

Chaitali Chakraborty, Kakali Bandyopadhyay, Shairee Ganguly, Bornini Banerjee and Shubham Mukherjee

Abstract

Fresh banana contains about 40% (w/w) peels. Banana peels are a good source of dietary fibres, polyphenols and some bioactive compounds. But in our society banana peels are considered as waste products. Even in banana processing industries banana peels are not utilized. As a waste material these peels can be hazardous to the environment as they can act as a medium of some harmful microorganisms. In the present study the nutraceutical properties of banana peels are utilized by using them in muffin production. Here banana peels are used as wheat flour substitute and it is added in different proportions (10%, 20%, and 30%) on the wet weight basis of the dough. The muffins were produced in microwave oven at different wattage (360W, 540W, 720W and 900W) as well as in baking oven at 110 °C temperature for 20 minutes. The polyphenol contents of all types of banana peel fortified muffins (10, 20 and 30% BPM) and control muffins, prepared in microwave oven and baking oven are determined in gm GAE/100 gm of samples by using five different extractions medium viz. acetone, methanol, ethanol, butanol and petroleum ether. It was observed that the polyphenols are mostly soluble in methanol for all types of BPMs as well as for control. For microwave processed 30% BPM the maximum polyphenols content is observed as 7.92 gm GAE/100 gm whereas this value is 7.77 gm GAE/100 gm for baking oven processed 30 % BPM in methanol. The sensory evaluations (by 9 point hedonic scale) of different BPMs are also carried out which showed the maximum overall acceptance for 30% BPM (microwave processed) and 20 % BPM (baking oven processed). Therefore raw banana peel can be utilized as a potent source of polyphenol in various foodstuffs.

Keywords: Banana peel muffin (BPM), polyphenol content, microwave processed, baking oven processed, sensory analysis

1. Introduction

The fruit and vegetable wastes (e.g. peels, seeds) are the non-product flows of raw materials whose economic values are less than the cost of collection and recovery for reuse; and therefore discarded as wastes. Phenolics are found in a plenty of plants and consist of an aromatic ring within the molecular structure (Singh *et al.* 2012) [10]. The agro-residues cannot be regarded as the wastes but become an additional valuable resource to augment existing natural materials. Recycling, reprocessing and eventual utilization of food processing residues offer potential of returning these by-products to beneficial uses rather than their discharge to the environment which cause detrimental environmental effects. Banana fruits contain various antioxidants such as gallic acid and dopamine. *Musa sapientum* which is commonly called banana is a herbaceous plant of the family *Musaceae*. Being as a tropical plant, banana protects itself from the oxidative stress caused by strong sunshine and high temperature by producing large amount of antioxidants. Interestingly, banana peel extracts have also been found to contain a high capacity to scavenge 2, 2-diphenyl-1-picrylhydrazyl (DPPH•) and 2,2'-azino-bis (3-ethylbenzothiazoline) -6-sulfonic acid (ABTS•+) free radicals (González-Montelongo *et al.*, 2010; Kedare and Singh, 2011) [4, 6]. Moreover, the extraction of antioxidants from banana peels is a great way for waste management because the main by-product from banana processing industry is its peel (Anal *et al.*, 2014) [2]. Sundaram *et al.*, 2011 [12], reported that the unripe banana peel sample had higher antioxidant potency than ripe and leaky ripe. Banana peel extract is classified as non-toxic to normal human cells criteria established by the National Cancer Standard Institute. (Anjum, *et al.* 2015), therefore, it can be safely utilized as a natural source of antioxidants and enzyme to cure disease.

Banana is a highly perishable and bulky fruit, which requires processing into a more stable and convenient form.



ISSN (E): 2277- 7695

ISSN (P): 2349-8242

NAAS Rating 2017: 5.03

TPI 2017; 6(10): 196-199

© 2017 TPI

www.thepharmajournal.com

Received: 14-08-2017

Accepted: 15-09-2017

Kakali Bandyopadhyay

Associate Professor, Department of Food Technology, Gurunanak Institute of Technology, 157/F, Nilgunj Road, Panihati, Sodepur, Kolkata, West Bengal, India

Chaitali Chakraborty

Associate Professor, Department of Dairy Chemistry, Faculty of Dairy Technology, WBUAFS, Mohanpur, Nadia, West Bengal, India

Suravi Chakraborty

Final Year B. Tech Student, Department of Food Technology, Gurunanak Institute of Technology, 157/F, Nilgunj Road, Panihati, Sodepur, Kolkata, West Bengal, India

Shailee Ganguly

Assistant Professor, Department of Food Technology, Gurunanak Institute of Technology, 157/F, Nilgunj Road, Panihati, Sodepur, Kolkata, West Bengal, India

Correspondence**Kakali Bandyopadhyay**

Associate Professor, Department of Food Technology, Gurunanak Institute of Technology, 157/F, Nilgunj Road, Panihati, Sodepur, Kolkata, West Bengal, India

Change in physicochemical properties of edible oil during frying: A review

Kakali Bandyopadhyay, Chaitali Chakraborty, Suravi Chakraborty and Shailee Ganguly

Abstract

Edible oils from plant, animal, or synthetic origin, are used in frying, baking, and other forms of cooking, and in salad dressings and bread dips. Plant-derived edible oils consist of carboxylic acids with long hydrocarbon chains, in contrast to petroleum-based oils which lack the carboxyl group on the end. The carboxyl group makes the oils edible, providing a site for human enzymes to attack and break down the chain in a process called beta-oxidation. There are a wide variety of cooking oils from plant sources such as olive oil, palm oil, soybean oil, canola oil (rapeseed oil), corn oil, peanut oil and other vegetable oils, as well as animal-based oils like butter and lard. This paper incorporates a comparative study of different types of frying oils by their physicochemical properties and compositional qualities. There are numerous health benefits of frying oils which has been covered in the paper along with the future of edible oils in India.

Keywords: Edible oils, carboxyl group, physicochemical properties, health benefits.

1. Introduction

Lipids and triacylglycerol naturally occur in oils and fats. Their chemical composition contains saturated and unsaturated fatty acids and glycerides. Edible oils are vital constituents of our daily diet, which provide energy, essential fatty acids and serve as a carrier of fat soluble vitamins (Erum Zahir *et al.*, 2014) [26]. Cooking oil is typically a liquid at room temperature, although some oils that contain saturated fat, such as coconut oil, palm oil and palm kernel oil are solid.

Fat frying is one of the oldest and popular food preparations. Fried foods have desirable flavour, colour and crispy texture, which make deep-fat fried foods very popular to consumers. Frying is a process of immersing food in hot oil with a contact among oil, air, and food at a high temperature of 150 to 190°C. The simultaneous heat and mass transfer of oil, food and air during fat frying produces the desirable and unique quality of fried foods. Frying oil acts as a heat transfer medium and contributes to the texture and flavour of fried food (Hassan A. Mudawi *et al.*, 2014) [15]. Numerous types of edible oils of plant and animal origin are used in frying, depending on regional availability. Palm oil is often used in Southeast Asia, coconut and groundnut oil in the Indian subcontinent, and olive oil in the Mediterranean region. During the last five decades, the Western food industry has become increasingly dependent on the frying process to manufacture a variety of snack foods. Fried foods such as potato chips, french fries, and fried fish and chicken have gained worldwide popularity (Farkas B.E. *et al.*, 1996) [10].

According to Erum Zahir, the quality of Corn and Mustard oils was analyzed by evaluating physicochemical properties such as density, viscosity, boiling point, peroxide, iodine and saponification values. Results are presented in Table 1. Oils with lower values of viscosity and density are highly appreciable to consumers. In order to design an advanced technological process these properties are very important parameters.

From the results obtained as presented in Table 1 the saponification value of palm kernel oil (280.5±56.1 mg KOH/g) is higher than those obtained for coconut oil (257.5±6.5 mg KOH/g) and groundnut oil (191.5±3.5 mg KOH/g) and since the higher the saponification value, the higher the unsaturated level of the oil, it can thus be inferred that palm kernel oil possess more unsaturated fatty acids than groundnut and coconut oils. It also indicates that the molecular weight of palm kernel oil is less than those of groundnut and coconut oils (Theodore, 1983) [23]. The iodine value obtained for palm kernel oil (i.e., 15.86±4.02 mgKOH/g) is also higher



ISSN (E): 2277- 7695
ISSN (P): 2349-8242
NAAS Rating 2017: 5.03
TPI 2017; 6(10): 421-424
© 2017 TPI
www.thepharmajournal.com
Received: 11-08-2017
Accepted: 12-09-2017

Chaitali Chakraorty

Associate Professor, Department of Dairy Chemistry, Faculty of Dairy Technology, WBUAFS, Mohanpur, Nadia, West Bengal, India

Kakali Bandyopadhyay

Associate Professor, Department of Food Technology, Guru Nanak Institute of Technology, Panihati, Sodepur, Kolkata, India

Riya Dasgupta

Final Year B.Tech Student, Department of Food Technology, Guru Nanak Institute of Technology, Panihati, Sodepur, Kolkata, India

Abhijit Ghosh

Final Year B.Tech Student, Department of Food Technology, Guru Nanak Institute of Technology, Panihati, Sodepur, Kolkata, India

Poulami Mitra

Final Year B.Tech Student, Department of Food Technology, Guru Nanak Institute of Technology, Panihati, Sodepur, Kolkata, India

Tania Paul

Final Year B.Tech Student, Department of Food Technology, Guru Nanak Institute of Technology, Panihati, Sodepur, Kolkata, India

Abhik Dutta

Final Year B.Tech Student, Department of Food Technology, Guru Nanak Institute of Technology, Panihati, Sodepur, Kolkata, India

Correspondence

Chaitali Chakraorty

Associate Professor, Department of Dairy Chemistry, Faculty of Dairy Technology, WBUAFS, Mohanpur, Nadia, West Bengal, India

Neutraceutical potential of some rare fruits & vegetables: A Review

Chaitali Chakraorty, Kakali Bandyopadhyay, Riya Dasgupta, Abhijit Ghosh, Poulami Mitra, Tania Paul and Abhik Dutta

Abstract

Vegetables are considered essential for well-balanced diets since they supply vitamins, minerals, dietary fiber, and phytochemicals. Each vegetable group contains a unique combination and amount of these phyto-nutraceuticals, which distinguishes them from other groups and vegetables within their own group. In the daily diet, rare vegetables can add more nutrition as well as play significant role in preventing various chronic diseases. Rare fruits and vegetables mainly include Broccoli, Purple Cabbage, Kiwi, Fig, Pomelo, Star fruit and many more. All of these are rich source of vitamins, minerals and carbohydrate with low fat content. Mainly they are preventive against different cancer and tumors that lead to cancer. The promotion of healthy rare vegetable products has coincided with a surging consumer interested in the healthy functionality of food.

Keywords: Rare fruits and vegetables, cancer, antioxidants, phytochemicals, and phyto-nutraceuticals

Introduction

The term vegetable refers to edible part(s) of a plant consumed raw or cooked, generally with a main dish, in a mixed dish, as an appetizer or in a salad. Vegetables include edible stems and stalks, roots, tubers, bulbs, leaves, flowers, some fruits, pulses (mature beans and peas), fungi (mushrooms, truffles), algae (seaweed) and sweet corn and hominy (cereal grains used as vegetables). Vegetables help to meet up major portion of the diet of humans through worldwide and play a significant role in human nutrition, especially as sources of phytonutraceuticals: vitamins (C, A, B1, B6, B9, E), minerals, dietary fiber and phytochemicals (Quebedeaux *et al.* 1990-1999) [18]. Some phytochemicals of vegetables act as strong antioxidants and plays significant role to reduce the risk of chronic disease by protecting against free-radical damage, by modifying metabolic activation and detoxification of carcinogens, or even influencing processes that alter the course of tumor cells (Southon *et al.*, 2000-2009) [23]. According to the 2007 World Health Report unbalanced diets with low vegetable intake and low consumption of complex carbohydrates and dietary fiber are estimated to cause some 2.7 million deaths each year, and were among the top 10 risk factors contributing to mortality (Dias, 2011) [5]. Especially, rare vegetables contributes significantly overall good health, improvement of gastrointestinal health and vision, reduced risk for some forms of cancer, heart disease, stroke, diabetes, anemia, gastric ulcer, rheumatoid arthritis, and other chronic diseases. Some rare vegetables are as follows: Broccoli, Purple Cabbage, Kiwi, Fig, Pomelo, Star fruit. Cruciferous vegetables (Brassicaceae or Cruciferae family) which include, cabbage, broccoli, cauliflower, Brussels sprouts, kales, kailan, chinese cabbage, turnip, rutabaga, radish, horseradish, rocket, watercress, mustards, among other vegetables, provide the richest sources of glucosinolates in the human diet.

Different Rare Vegetables

1. Broccoli: *Broccoli* ("*Brassica oleracea var. italic*") is a Cruciferous green leaf Cole vegetable. This plant is native of Italy, but can be successfully grown in India also. It is a rich source of valuable nutrients Vitamin A, C & riboflavin. It contains high amount of Iron and Calcium and is a non-fattening food and possesses various medicinal properties as well (Mishra and Mukherjee, 2012) [15]. Broccoli grows best in temperature ranging between 18 °C and 23° C. Sprouting broccoli has a larger number of heads with many thin stalks. It is planted in May to be harvested during the winter in areas with temperate climates. Romanesco broccoli has a distinctive fractal appearance of its heads, and is yellow-green in color. It is technically the *Botrytis* (Cauliflower) cultivar group. Purple cauliflower is a type of broccoli



ISSN (E): 2277- 7695

ISSN (P): 2349-8242

NAAS Rating 2017: 5.03

TPI 2017; 6(11): 93-101

© 2017 TPI

www.thepharmajournal.com

Received: 15-09-2017

Accepted: 16-10-2017

Chaitali Chakraborty

Associate Professor, Department of Dairy Chemistry, Faculty of Dairy Technology, WBUAFS, Mohanpur, Nadia, West Bengal, India

Kakali Bandyopadhyay

Associate Professor, Department of Food Technology, Guru Nanak Institute of Technology, Panihati, Sodepur, Kolkata, West Bengal, India

Chandralekha Bhowmik

Final Year B.Tech Student, Department of Food Technology, Guru Nanak Institute of Technology, Panihati, Sodepur, Kolkata, West Bengal, India

Prostuti Chakravorty

Final Year B.Tech Student, Department of Food Technology, Guru Nanak Institute of Technology, Panihati, Sodepur, Kolkata, West Bengal, India

Rupsa Roychowdhury

Final Year B.Tech Student, Department of Food Technology, Guru Nanak Institute of Technology, Panihati, Sodepur, Kolkata, West Bengal, India

Shubhapriya Samanta

Final Year B.Tech Student, Department of Food Technology, Guru Nanak Institute of Technology, Panihati, Sodepur, Kolkata, West Bengal, India

Manasi Roy

Final Year B.Tech Student, Department of Food Technology, Guru Nanak Institute of Technology, Panihati, Sodepur, Kolkata, West Bengal, India

Correspondence**Chaitali Chakraborty**

Associate Professor, Department of Dairy Chemistry, Faculty of Dairy Technology, WBUAFS, Mohanpur, Nadia, West Bengal, India

Utilization of various seeds: A review

Chaitali Chakraborty, Kakali Bandyopadhyay, Chandralekha Bhowmik, Prostuti Chakravorty, Rupsa Roychowdhury, Shubhapriya Samanta and Manasi Roy

Abstract

Seed, known to be an embryo possessing part of a plant, enclosed in a protective outer covering, does not have much utilization known to us. Seeds which are generally thrown away as a waste product have various benefits hidden among themselves. In this review paper, seeds and their different utilizations have been highlighted. We see that seeds, because of their concealed polyphenols and antimicrobial activities, serve us with various medicinal purposes. By-products such as seed flour and seed oil derived from them are enriched with nutrition value and antioxidant properties. Thus, instead of being wasted and withered, proper utilization of seeds should be done.

Keywords: embryo, polyphenols, antimicrobial activities, seed flour, seed oil

1. Introduction

Seed has been considered as a waste product dumped from households or fruit/vegetable processing industries. But seeds have found their utilization in food, pharmaceutical and cosmetic industries for their antioxidant and medicinal values. Besides, seed can be used for processing of seed flour, seed oil and can be utilized as a source of various polyphenols and flavonoids. The parts of the plant most commonly used for the therapeutic purposes in the "Alternative Medicinal" systems are the seeds which are contained in an inflated capsule formed from the united follicles containing considerable amount of oil having pungent and bitter taste. (Padhye *et al.* 2008). (Umbelliferae). Essential oil of fennel is used as flavoring agents in food products such as beverages, bread, pickles, pastries, and cheese. It is also used as a constituent of cosmetic and pharmaceutical products (Piccaglia *et al.*, 2001) [75]. Herbal drugs and essential oils of fennel have hepatoprotective effects (Ozbek *et al.* 2003) [71], as well as antispasmodic effects. They are also known for their diuretic, anti-inflammatory, analgesic and antioxidant activities (Choi, E, 2004) [31] (Anand *et al.*, 2008) [5] reported that fennel seed possesses anticancer activity.

In general, the use of food parts usually discarded by industries adds nutrients to various preparations. (Storck *et al.* 2013) [84-89], elaborated preparations using papaya seed cake and papaya peel jam and observed an increased fiber content and sensory analyses were conducted. *Sesamum indicum* L. (Pedaliaceae) is an annual shrub with white bell-shaped flowers with a hint of blue, red or yellow with branches or without branches. It is grown for the production of seeds which is rich in oil content. (Chakraborty *et al.* 2008). Commonly the seeds are used primarily as a spice and food preservative. In folk medicinal practices they are ingested with food or mixed with honey and are primarily used as lactogogues, carminative and antihelmthic agents. The seeds have also been used as diuretics, anti-hypertensive, muscle relaxants and as immunity enhancers in immune-compromised people. Importantly, the seeds have been reported to be safe when used orally in moderate amount in food (Der Marderosian. *et al.*, 2005) [34]. Several beneficial pharmacological effects have been attributed to various crude or purified components of these seeds including antihistaminic (Chakravorty, 1993) [30], antihypertensive (Zaoui *et al.*, 2000) [98], hypoglycemic (Al-Hader *et al.*, 1993) [1], antifungal (Khan *et al.*, 2003) [51], anti-inflammatory (Al-Ghamdi, 2001) [9] along with significant anti-neoplastic (Worthen *et al.*, 1998) [96] activities. These studies collectively provide early indication that further development of agents derived from black cumin seeds could be useful in modern medicine.

The chemical composition of fenugreek seed (FS) has been thoroughly studied and its medicinal properties are associated with its phytochemicals such as galactomannans, phenolic compounds, alkaloids, proteins, vitamins (A, B1, C and nicotinic acid) and volatile oils

Original Research Article

<https://doi.org/10.20546/ijcmas.2018.705.167>

**A Study on Antimicrobial Properties and Medicinal Value of
Adhatoda vasica, *Centella asiatica*, *Paederia foetida*,
Nyctanthes arbor-tristis, *Ocimum tenuiflorum***

Anju Paul^{1*}, Amit Kumar Barman², Pradip Kumar Roy² and Subhajit Ray³

¹Department of Food Technology, Guru Nanak Institute of Technology, India

²Faculty of Dairy Technology, WBUAFS, Mohanpur Campus, W.B., Nadia-741252, India

³Department of Food Engineering and Technology, CIT, Assam-783370, India

**Corresponding author*

ABSTRACT

Keywords

Medicinal plants,
Pharmacological
activity, herbs,
polyphenols

Article Info

Accepted:

12 April 2018

Available Online:

10 May 2018

The use of various herbal remedies and preparations are described throughout human history representing the origin of modern medicine. Herbal medicine is also called botanical medicine or phyto-medicine, and is defined as the use whole plants or part of plants to prevent or treat illness. Medicinal plants constitute an important component of flora and are widely distributed in India. The pharmacological evaluation of substances from plants is an established method for the identification of lead compounds which can leads to the development of novel and safe medicinal agents. The importance of medicinal plants and traditional health systems in solving the health care problems of the world is gaining increasing attention. Because of this resurgence of interest, the research on plants of medicinal importance is growing phenomenally at the international level, often to the detriment of natural habitats and mother populations in the countries of origin. The study on antimicrobial and medicinal values of: *Adhatoda vasica* (*vasaka*), *Centella asiatica* (*thankuni*), *Paederia foetida* (*gadal*), *Nyctanthes arbor-tristis* (*shiuli*), will help us to understand their anti-microbial property so that they can be used in more effective and efficient manner for pharmacological purposes.

Introduction

World Health Organisation (WHO) has defined medicinal plants as plants that contain properties or compounds that can be used for therapeutic purposes or those that synthesize metabolites to produce useful drugs. Medicinal plants constitute an important component of flora and are widely distributed in India. The pharmacological evaluation of substances from plants is an established

method for the identification of lead compounds which can leads to the development of novel and safe medicinal agents. The importance of medicinal plants and traditional health systems in solving the health care problems of the world is gaining increasing attention. Because of this resurgence of interest, the research on plants of medicinal importance is growing phenomenally at the international level, often to the detriment of natural habitats and mother

CLIMATIC CONDITIONS REQUIRED AND FACTORS AFFECTING THE PRODUCTION OF SWEET POTATO

CHAITALI CHAKRABORTY¹, RIYA DASGUPTA² & DEBJIT GHOSH³

^{1,2}Department of Food Technology, Guru Nanak Institute of Technology, Panihati, Sodepur, Kolkata, India

³Don-Limon GmbH, Hamburg, Germany

ABSTRACT:

Sweet potato is a subtropical crop. It requires moderate temperature ranging between 21-26°C. It requires the average temperature 75°F (24°C) for growth. The other factor that also have significant effect on growth are abundant sunlight, warm nights, annual rainfall [(750-1000mm), with minimum range of 500mm in growing time]. Climate change such as global warming, elevated CO₂ concentration, high temperature etc affects sweet potato production worldwide. Elevated CO₂ cause malformation of tuber. High temperature also causes reduced yield. Overall climate change affects sweet potato greatly.

KEYWORDS: Climate Change, Production, Yield, Carbon-di-Oxide & Temperature

Received: Dec 08, 2016; **Accepted:** Jan 06, 2017; **Published:** Feb 04, 2017; **Paper Id.:** IJASRFEB201758

INTRODUCTION

Sweet potatoes can be grown where there is a long frost-free period with warm temperatures in the growing season. The plant does not tolerate frost. Sweet potato requires moderate temperature (21-26°C, so, it can be grown in every month but in different countries worldwide. Sweet potatoes require plenty of sunshine, but shade causes yield reduction. However, sweet potato is intercropped with other seasonal crops like pigeon pea, maize, etc (Nedunchezhiyan *et al.*, 2010) and to fulfill the objective of intensification of crop and to gain the profit at a maximum level, it can be grown as an intercrop. At initiation stage (0-60) days, sweet potato is very much sensitive to draught and water-logging. If it is not maintained properly, then it causes rotting of tuber and reduced growth of storage root. (Nedunchezhiyan and Ray 2010). Sweet potato is very much affected by saline and alkaline condition of the soil. (Dasgupta *et al.* 2006, Mukherjee *et al.* 2006).

FACTORS AFFECTING THE PRODUCTION OF SWEET POTATO

Concentration of CO₂

Increased rates of photosynthesis, WUE, and NUE on plants grown at elevated CO₂ concentrations not only alter biomass, but also change the plants elemental composition. The majority of studies looking at these effects have focused on the foliar portion of crops, or the seed or grain of major global crops and resulted in increased carbohydrate concentration and decreased protein and mineral concentration (Seneweer and Conroy, 1997; Fangmeier *et al.*, 1999; Prior *et al.*, 1998). Loladze (2002) reviewed 25 studies that looked at changes in mineral concentration of plants grown under slightly elevated (twice ambient) CO₂ concentrations. He found average percent decreases in N, P, K, Ca, S, Mg, Fe, Zn, Mn, and Cu for all foliar plants, and decreases for all minerals except K (.86% increase) and Cu (no data) for wheat showing the same trends found for wheat mineral

Development Of Papaya Peel Flour Based Cookies And Evaluation Of Its Quality

Dr. Kanta Bokaria

Post-Doctoral Fellow, Department of Food
Technology, Guru Nanak Institute of
Technology, 157/F Nilgunj Road, Sodepur,
Panihati, Kolkata-700114, West Bengal, India.
Email- kantabokaria@gmail.com

Dr. Subhajit Ray

Associate Professor &HOD, Department of Food
Technology, Guru Nanak Institute of Technology,
157/F Nilgunj Road, Sodepur, Panihati, Kolkata-
700114, West Bengal, India. Email-
head_ft.gnit@jisgroup.org

Abstract—The focus of this research project was to develop papaya peel flour from raw papaya and utilizing the papaya peel flour for the development of value added cookies. Fresh papaya was blanched, peeled, cut into small pieces, further washed and treated with sanitizer, tray dried and finally grinded to powder form. Prepared papaya peel flour was evaluated for chemical analysis. In the present study the moisture (%), ash (%), acidity, fat (%), insoluble solid(%),soluble solid(%), protein content(%), vitamin C, total fiber, total carbohydrate and antioxidant content of papaya peel flour were estimated as 13.63%, 5.25%, .163% ,.1%, 80%, 20%, 8.64%,.221.U, 33.5%, 38.88% and 514.6mg/100gm respectively. Papaya peel flour fortified biscuits were formulated by incorporating 5%, 7.5% and 10% papaya peel flour with ordinary flour in the present investigation. Experimental result revealed 5% peel flour formulated cookies gives the best results in terms of physicochemical quality. The physicochemical and nutritional quality parameters viz. moisture(%), ash(%), acidity, fat(%),vitamin C, insoluble solid (%),soluble solid(%),protein content(%),total fiber, total carbohydrate and antioxidant content of all fortified varieties of cookies were determined. The best result obtained for 5% papaya peel fortified cookies and was determined as 10.35%, 4.5%, 0.02%, 15.4%, .31U, 85%, 15%,9.3%,.017%,60.43% and 9gm/100gm respectively. Now the experimental result revealed that papaya peel flour fortified cookies contains significant amount of protein and antioxidant in comparison to it's ordinary counterpart which improves it's nutritional characteristics. Sensory evaluations of all fortified varieties were also carried out and it was found that 5% papaya peel flour blend for cookies formulation was acceptable. Therefore 5% papaya peel flour based formulated cookies proves enhanced nutritional properties, physicochemical characteristics and organoleptic attributes.

Keywords—*Papaya peel flour, value addition, antioxidant, sensory evaluation, nutrition*

I. INTRODUCTION

Papaya (*Carica papaya L.*) belongs to the family Caricaceae. Papaya (*C. papaya L.*) is the fourth most

important tropical fruit around the globe (Scheldeman et al. 2007). The major producers of papaya in the world are Australia, United States, Philippines, SriLanka, South Africa, India, Bangladesh, Malaysia and a number of other countries in tropical America (Anuara et al. 2008). Some of the active compounds in papaya are ascorbic acid (108 mg/100 g), antioxidant, β -carotene, α -tocopherol, flavonoids, vitamin B1, papain and niacin (Leontowicz et al. 2007; Lim et al. 2007). Fruit processing results in large amounts of waste as by products such as peels and seeds. Thus, by-products and its further utilization in the production of food additives or supplements with high nutritional value have aroused great interest since they are high-value products. It is very much interesting that such wastes are important sources of sugars, minerals, organic acids, fiber, and phenolic compounds that have a wide range of pharmacological activities, which include antitumor, antiviral, antibacterial, cardio protective, and ant mutagenic activities (Djilas et al., 2009). In addition, these components help to maintain health and prevent diseases such as cancer, cardiovascular and many other degenerative diseases (Larrauri 1999). Fortification of these nutritional constituents in processed food products is a way to increase daily cooking by creating new recipes such as jellies, pies, juice, and pastries, in addition to nutritionally enriched diets, providing more fiber, vitamins, and minerals (Storck et al., 2013). (Storck et al., 2013) elaborated preparations using papaya seed cake and papaya peel jam and observed an increase in fiber content and sensory analysis were conducted. In many countries, cookies are generally consumed and it could be considered as a vehicle for nutrient transfer (Arshad et al. 2007). A number of studies have been reported on the improved nutritive value of high protein cookies made of composite flour such as a blend of soybean (Shrestha and Noomhorm 2002) and unripe banana flour replacing the wheat flour. Rice and black bean extruded flour replacing wheat flour can be considered as a good source of protein and fiber and reducing lipids in cookies (Lund and Smoot 1982). This study was conducted with the objective to produce and analyze papaya waste (peel) flour for its chemical constituents with the purpose to use it in preparations and formulations of nutritionally enriched value added cookies in terms of nutritional properties, physicochemical characteristics and organoleptic attributes.